

# CHAPTER 2

## Differentiation

### Section 2.1 The Derivative and the Tangent Line Problem


1. At  $(x_1, y_1)$ , slope = 0.

At  $(x_2, y_2)$ , slope =  $\frac{5}{2}$ .

2. At  $(x_1, y_1)$ , slope =  $\frac{2}{3}$ .

At  $(x_2, y_2)$ , slope =  $-\frac{2}{5}$ .

3. (a), (b)



(c)  $y = \frac{f(4) - f(1)}{4 - 1}(x - 1) + f(1)$

$$= \frac{3}{3}(x - 1) + 2$$

$$= 1(x - 1) + 2$$

$$= x + 1$$

4. (a)  $\frac{f(4) - f(1)}{4 - 1} = \frac{5 - 2}{3} = 1$

$$\frac{f(4) - f(3)}{4 - 3} \approx \frac{5 - 4.75}{1} = 0.25$$

So,  $\frac{f(4) - f(1)}{4 - 1} > \frac{f(4) - f(3)}{4 - 3}$ .

(b) The slope of the tangent line at  $(1, 2)$  equals  $f'(1)$ .

This slope is steeper than the slope of the line

through  $(1, 2)$  and  $(4, 5)$ . So,  $\frac{f(4) - f(1)}{4 - 1} < f'(1)$ .

5.  $f(x) = 3 - 5x$  is a line. Slope =  $-5$

6.  $g(x) = \frac{3}{2}x + 1$  is a line. Slope =  $\frac{3}{2}$

7. Slope at  $(2, -5) = \lim_{\Delta x \rightarrow 0} \frac{g(2 + \Delta x) - g(2)}{\Delta x}$

$$= \lim_{\Delta x \rightarrow 0} \frac{(2 + \Delta x)^2 - 9 - (-5)}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{4 + 4(\Delta x) + (\Delta x)^2 - 4}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} (4 + \Delta x) = 4$$

8. Slope at  $(3, -4) = \lim_{\Delta x \rightarrow 0} \frac{f(3 + \Delta x) - f(3)}{\Delta x}$

$$= \lim_{\Delta x \rightarrow 0} \frac{5 - (3 + \Delta x)^2 - (-4)}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{5 - 9 - 6(\Delta x) - (\Delta x)^2 + 4}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{-6(\Delta x) - (\Delta x)^2}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} (-6 - \Delta x) = -6$$

9. Slope at  $(0, 0) = \lim_{\Delta t \rightarrow 0} \frac{f(0 + \Delta t) - f(0)}{\Delta t}$

$$= \lim_{\Delta t \rightarrow 0} \frac{3(\Delta t) - (\Delta t)^2 - 0}{\Delta t}$$

$$= \lim_{\Delta t \rightarrow 0} (3 - \Delta t) = 3$$

10. Slope at  $(1, 5) = \lim_{\Delta t \rightarrow 0} \frac{h(1 + \Delta t) - h(1)}{\Delta t}$

$$= \lim_{\Delta t \rightarrow 0} \frac{(1 + \Delta t)^2 + 4(1 + \Delta t) - 5}{\Delta t}$$

$$= \lim_{\Delta t \rightarrow 0} \frac{1 + 2(\Delta t) + (\Delta t)^2 + 4 + 4(\Delta t) - 5}{\Delta t}$$

$$= \lim_{\Delta t \rightarrow 0} \frac{6(\Delta t) + (\Delta t)^2}{\Delta t}$$

$$= \lim_{\Delta t \rightarrow 0} (6 + \Delta t) = 6$$

11.  $f(x) = 7$

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{7 - 7}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} 0 = 0$$

12.  $g(x) = -3$

$$g'(x) = \lim_{\Delta x \rightarrow 0} \frac{g(x + \Delta x) - g(x)}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{-3 - (-3)}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{0}{\Delta x} = 0$$

13.  $f(x) = -10x$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{-10(x + \Delta x) - (-10x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{-10x - 10\Delta x + 10x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{-10\Delta x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (-10) = -10 \end{aligned}$$

14.  $f(x) = 7x - 3$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{7(x + \Delta x) - 3 - (7x - 3)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{7x + 7\Delta x - 3 - 7x + 3}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{7(\Delta x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} 7 = 7 \end{aligned}$$

17.  $f(x) = x^2 + x - 3$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^2 + (x + \Delta x) - 3 - (x^2 + x - 3)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^2 + 2x(\Delta x) + (\Delta x)^2 + x + \Delta x - 3 - x^2 - x + 3}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{2x(\Delta x) + (\Delta x)^2 + \Delta x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (2x + \Delta x + 1) = 2x + 1 \end{aligned}$$

18.  $f(x) = x^2 - 5$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^2 - 5 - (x^2 - 5)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^2 + 2x(\Delta x) + (\Delta x)^2 - 5 - x^2 + 5}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{2x(\Delta x) + (\Delta x)^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (2x + \Delta x) = 2x \end{aligned}$$

15.  $h(s) = 3 + \frac{2}{3}s$

$$\begin{aligned} h'(s) &= \lim_{\Delta s \rightarrow 0} \frac{h(s + \Delta s) - h(s)}{\Delta s} \\ &= \lim_{\Delta s \rightarrow 0} \frac{3 + \frac{2}{3}(s + \Delta s) - \left(3 + \frac{2}{3}s\right)}{\Delta s} \\ &= \lim_{\Delta s \rightarrow 0} \frac{3 + \frac{2}{3}s + \frac{2}{3}\Delta s - 3 - \frac{2}{3}s}{\Delta s} \\ &= \lim_{\Delta s \rightarrow 0} \frac{\frac{2}{3}\Delta s}{\Delta s} = \frac{2}{3} \end{aligned}$$

16.  $f(x) = 5 - \frac{2}{3}x$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{5 - \frac{2}{3}(x + \Delta x) - \left(5 - \frac{2}{3}x\right)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{5 - \frac{2}{3}x - \frac{2}{3}\Delta x - 5 + \frac{2}{3}x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{-\frac{2}{3}(\Delta x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \left(-\frac{2}{3}\right) = -\frac{2}{3} \end{aligned}$$

19.  $f(x) = x^3 - 12x$

$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{[(x + \Delta x)^3 - 12(x + \Delta x)] - [x^3 - 12x]}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{x^3 + 3x^2\Delta x + 3x(\Delta x)^2 + (\Delta x)^3 - 12x - 12\Delta x - x^3 + 12x}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{3x^2\Delta x + 3x(\Delta x)^2 + (\Delta x)^3 - 12\Delta x}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} (3x^2 + 3x\Delta x + (\Delta x)^2 - 12) = 3x^2 - 12
 \end{aligned}$$

20.  $f(x) = x^3 + x^2$

$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{[(x + \Delta x)^3 + (x + \Delta x)^2] - [x^3 + x^2]}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{x^3 + 3x^2\Delta x + 3x(\Delta x)^2 + (\Delta x)^3 + x^2 + 2x\Delta x + (\Delta x)^2 - x^3 - x^2}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{3x^2\Delta x + 3x(\Delta x)^2 + (\Delta x)^3 + 2x\Delta x + (\Delta x)^2}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} (3x^2 + 3x\Delta x + (\Delta x)^2 + 2x + (\Delta x)) = 3x^2 + 2x
 \end{aligned}$$

21.  $f(x) = \frac{1}{x-1}$

$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{\frac{1}{x + \Delta x - 1} - \frac{1}{x - 1}}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{(x - 1) - (x + \Delta x - 1)}{\Delta x(x + \Delta x - 1)(x - 1)} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{-\Delta x}{\Delta x(x + \Delta x - 1)(x - 1)} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{-1}{(x + \Delta x - 1)(x - 1)} \\
 &= -\frac{1}{(x - 1)^2}
 \end{aligned}$$

22.  $f(x) = \frac{1}{x^2}$

$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{\frac{1}{(x + \Delta x)^2} - \frac{1}{x^2}}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{x^2 - (x + \Delta x)^2}{\Delta x(x + \Delta x)^2 x^2} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{-2x\Delta x - (\Delta x)^2}{\Delta x(x + \Delta x)^2 x^2} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{-2x - \Delta x}{(x + \Delta x)^2 x^2} \\
 &= \frac{-2x}{x^4} \\
 &= -\frac{2}{x^3}
 \end{aligned}$$

$$23. f(x) = \sqrt{x+4}$$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x + \Delta x + 4} - \sqrt{x + 4}}{\Delta x} \cdot \left( \frac{\sqrt{x + \Delta x + 4} + \sqrt{x + 4}}{\sqrt{x + \Delta x + 4} + \sqrt{x + 4}} \right) \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x + 4) - (x + 4)}{\Delta x [\sqrt{x + \Delta x + 4} + \sqrt{x + 4}]} \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{\sqrt{x + \Delta x + 4} + \sqrt{x + 4}} = \frac{1}{\sqrt{x + 4} + \sqrt{x + 4}} = \frac{1}{2\sqrt{x + 4}} \end{aligned}$$

$$24. f(x) = \frac{4}{\sqrt{x}}$$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\frac{4}{\sqrt{x + \Delta x}} - \frac{4}{\sqrt{x}}}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{4\sqrt{x} - 4\sqrt{x + \Delta x}}{\Delta x \sqrt{x} \sqrt{x + \Delta x}} \cdot \left( \frac{\sqrt{x} + \sqrt{x + \Delta x}}{\sqrt{x} + \sqrt{x + \Delta x}} \right) \\ &= \lim_{\Delta x \rightarrow 0} \frac{4x - 4(x + \Delta x)}{\Delta x \sqrt{x} \sqrt{x + \Delta x} (\sqrt{x} + \sqrt{x + \Delta x})} \\ &= \lim_{\Delta x \rightarrow 0} \frac{-4}{\sqrt{x} \sqrt{x + \Delta x} (\sqrt{x} + \sqrt{x + \Delta x})} \\ &= \frac{-4}{\sqrt{x} \sqrt{x} (\sqrt{x} + \sqrt{x})} = \frac{-2}{x\sqrt{x}} \end{aligned}$$

$$25. (a) f(x) = x^2 + 3$$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{[(x + \Delta x)^2 + 3] - (x^2 + 3)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^2 + 2x\Delta x + (\Delta x)^2 + 3 - x^2 - 3}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{2x\Delta x + (\Delta x)^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (2x + \Delta x) = 2x \end{aligned}$$

At  $(-1, 4)$ , the slope of the tangent line is

$$m = 2(-1) = -2.$$


The equation of the tangent line is

$$y - 4 = -2(x + 1)$$

$$y - 4 = -2x - 2$$

$$y = -2x + 2$$

(b)



(c) Graphing utility confirms  $\frac{dy}{dx} = -2$  at  $(-1, 4)$ .

26. (a)  $f(x) = x^2 + 2x - 1$

$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{[(x + \Delta x)^2 + 2(x + \Delta x) - 1] - [x^2 + 2x - 1]}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{[x^2 + 2x\Delta x + (\Delta x)^2 + 2x + 2\Delta x - 1] - [x^2 + 2x - 1]}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{2x\Delta x + (\Delta x)^2 + 2\Delta x}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} (2x + \Delta x + 2) = 2x + 2
 \end{aligned}$$

At  $(1, 2)$ , the slope of the tangent line is  $m = 2(1) + 2 = 4$ .


The equation of the tangent line is

$$y - 2 = 4(x - 1)$$

$$y - 2 = 4x - 4$$

$$y = 4x - 2.$$

(b)



(c) Graphing utility confirms  $\frac{dy}{dx} = 4$  at  $(1, 2)$ .

27. (a)  $f(x) = x^3$

$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^3 - x^3}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{x^3 + 3x^2\Delta x + 3x(\Delta x)^2 + (\Delta x)^3}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{3x^2\Delta x + 3x(\Delta x)^2 + (\Delta x)^3}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} (3x^2 + 3x\Delta x + (\Delta x)^2) = 3x^2
 \end{aligned}$$


At  $(2, 8)$ , the slope of the tangent is  $m = 3(2)^2 = 12$ . The equation of the tangent line is

$$y - 8 = 12(x - 2)$$

$$y - 8 = 12x - 24$$

$$y = 12x - 16.$$

(b)




(c) Graphing utility confirms  $\frac{dy}{dx} = 12$  at  $(2, 8)$ .

28. (a)  $f(x) = x^3 + 1$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{[(x + \Delta x)^3 + 1] - (x^3 + 1)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^3 + 3x^2(\Delta x) + 3x(\Delta x)^2 + (\Delta x)^3 + 1 - x^3 - 1}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} [3x^2 + 3x(\Delta x) + (\Delta x)^2] = 3x^2 \end{aligned}$$

At  $(-1, 0)$ , the slope of the tangent line is  $m = 3(-1)^2 = 3$ . The equation of the tangent line is

$$\begin{aligned} y - 0 &= 3(x + 1) \\ y &= 3x + 3. \end{aligned}$$



(c) Graphing utility confirms  $\frac{dy}{dx} = 3$  at  $(-1, 0)$ .


29. (a)  $f(x) = \sqrt{x}$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x + \Delta x} - \sqrt{x}}{\Delta x} \cdot \frac{\sqrt{x + \Delta x} + \sqrt{x}}{\sqrt{x + \Delta x} + \sqrt{x}} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x) - x}{\Delta x(\sqrt{x + \Delta x} + \sqrt{x})} \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{\sqrt{x + \Delta x} + \sqrt{x}} = \frac{1}{2\sqrt{x}} \end{aligned}$$

At  $(1, 1)$ , the slope of the tangent line is  $m = \frac{1}{2\sqrt{1}} = \frac{1}{2}$ .

The equation of the tangent line is

$$\begin{aligned} y - 1 &= \frac{1}{2}(x - 1) \\ y - 1 &= \frac{1}{2}x - \frac{1}{2} \\ y &= \frac{1}{2}x + \frac{1}{2}. \end{aligned}$$



(c) Graphing utility confirms  $\frac{dy}{dx} = \frac{1}{2}$  at  $(1, 1)$ .

30. (a)  $f(x) = \sqrt{x-1}$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x + \Delta x - 1} - \sqrt{x - 1}}{\Delta x} \cdot \left( \frac{\sqrt{x + \Delta x - 1} + \sqrt{x - 1}}{\sqrt{x + \Delta x - 1} + \sqrt{x - 1}} \right) \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x - 1) - (x - 1)}{\Delta x(\sqrt{x + \Delta x - 1} + \sqrt{x - 1})} \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{\sqrt{x + \Delta x - 1} + \sqrt{x - 1}} = \frac{1}{2\sqrt{x-1}} \end{aligned}$$

At (5, 2), the slope of the tangent line is  $m = \frac{1}{2\sqrt{5-1}} = \frac{1}{4}$ .

The equation of the tangent line is

$$y - 2 = \frac{1}{4}(x - 5)$$

$$y - 2 = \frac{1}{4}x - \frac{5}{4}$$

$$y = \frac{1}{4}x + \frac{3}{4}$$

31. (a)  $f(x) = x + \frac{4}{x}$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x) + \frac{4}{x + \Delta x} - \left(x + \frac{4}{x}\right)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x(x + \Delta x)(x + \Delta x) + 4x - x^2(x + \Delta x) - 4(x + \Delta x)}{x(\Delta x)(x + \Delta x)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^3 + 2x^2(\Delta x) + x(\Delta x)^2 - x^3 - x^2(\Delta x) - 4(\Delta x)}{x(\Delta x)(x + \Delta x)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^2(\Delta x) + x(\Delta x)^2 - 4(\Delta x)}{x(\Delta x)(x + \Delta x)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^2 + x(\Delta x) - 4}{x(x + \Delta x)} \\ &= \frac{x^2 - 4}{x^2} = 1 - \frac{4}{x^2} \end{aligned}$$


At (-4, -5), the slope of the tangent line is  $m = 1 - \frac{4}{(-4)^2} = \frac{3}{4}$ .

The equation of the tangent line is

$$y + 5 = \frac{3}{4}(x + 4)$$


$$y + 5 = \frac{3}{4}x + 3$$

$$y = \frac{3}{4}x - 2$$



(c) Graphing utility confirms

$$\frac{dy}{dx} = \frac{1}{4} \text{ at } (5, 2).$$



(c) Graphing utility confirms

$$\frac{dy}{dx} = \frac{3}{4} \text{ at } (-4, -5).$$

$$32. (a) f(x) = x + \frac{6}{x+2}$$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\frac{6}{(x + \Delta x) + 2} - \frac{6}{x + 2}}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{6x + 12 - 6(x + \Delta x + 2)}{\Delta x(x + \Delta x + 2)(x + 2)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{6x + 12 - 6x - 6\Delta x - 12}{\Delta x(x + \Delta x + 2)(x + 2)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{-6\Delta x}{\Delta x(x + \Delta x + 2)(x + 2)} \\ &= \frac{-6}{(x + 2)^2} \end{aligned}$$

At  $(0, 3)$ , the slope of the tangent line is

$$m = -\frac{6}{4} = -\frac{3}{2}.$$


The equation of the tangent line is

$$y - 3 = -\frac{3}{2}(x - 0)$$

$$y - 3 = -\frac{3}{2}x$$

$$y = -\frac{3}{2}x + 3.$$

(b)



(c) Graphing utility confirms  $\frac{dy}{dx} = -\frac{3}{2}$  at  $(0, 3)$ .

33. Using the limit definition of derivative,  $f'(x) = 2x$ .

Because the slope of the given line is 2, you have

$$2x = 2$$

$$x = 1$$

At the point  $(1, 1)$  the tangent line is parallel to

$2x - y + 1 = 0$ . The equation of this line is

$$y - 1 = 2(x - 1)$$

$$y = 2x - 1.$$

34. Using the limit definition of derivative,  $f'(x) = 4x$ .

Because the slope of the given line is  $-4$ , you have

$$4x = -4$$

$$x = -1.$$

At the point  $(-1, 2)$  the tangent line is parallel to

$4x + y + 3 = 0$ . The equation of this line is

$$y - 2 = -4(x + 1)$$

$$y = -4x - 2.$$

35. From Exercise 27 we know that  $f'(x) = 3x^2$ .

Because the slope of the given line is 3, you have

$$3x^2 = 3$$

$$x = \pm 1.$$

Therefore, at the points  $(1, 1)$  and  $(-1, -1)$  the tangent

lines are parallel to  $3x - y + 1 = 0$ .

These lines have equations

$$y - 1 = 3(x - 1) \quad \text{and} \quad y + 1 = 3(x + 1)$$

$$y = 3x - 2 \quad \quad \quad y = 3x + 2.$$

36. Using the limit definition of derivative,  $f'(x) = 3x^2$ .

Because the slope of the given line is 3, you have

$$3x^2 = 3$$

$$x^2 = 1 \Rightarrow x = \pm 1.$$

Therefore, at the points  $(1, 3)$  and  $(-1, 1)$  the tangent

lines are parallel to  $3x - y - 4 = 0$ . These lines have equations

$$y - 3 = 3(x - 1) \quad \text{and} \quad y - 1 = 3(x + 1)$$

$$y = 3x \quad \quad \quad y = 3x + 4.$$

37. Using the limit definition of derivative,

$$f'(x) = \frac{-1}{2x\sqrt{x}}.$$

Because the slope of the given line is  $-\frac{1}{2}$ , you have

$$-\frac{1}{2x\sqrt{x}} = -\frac{1}{2}$$

$$x = 1.$$

Therefore, at the point  $(1, 1)$  the tangent line is parallel to

$x + 2y - 6 = 0$ . The equation of this line is

$$y - 1 = -\frac{1}{2}(x - 1)$$

$$y - 1 = -\frac{1}{2}x + \frac{1}{2}$$

$$y = -\frac{1}{2}x + \frac{3}{2}.$$



38. Using the limit definition of derivative,

$$f'(x) = \frac{-1}{2(x-1)^{3/2}}$$

Because the slope of the given line is  $-\frac{1}{2}$ , you have

$$\frac{-1}{2(x-1)^{3/2}} = -\frac{1}{2}$$

$$1 = (x-1)^{3/2}$$


$$1 = x-1 \Rightarrow x = 2.$$

At the point  $(2, 1)$ , the tangent line is parallel to  $x + 2y + 7 = 0$ . The equation of the tangent line is


$$y - 1 = -\frac{1}{2}(x - 2)$$

$$y = -\frac{1}{2}x + 2.$$


39. The slope of the graph of  $f$  is 1 for all  $x$ -values.




40. The slope of the graph of  $f$  is 0 for all  $x$ -values.




41. The slope of the graph of  $f$  is negative for  $x < 4$ , positive for  $x > 4$ , and 0 at  $x = 4$ .




42. The slope of the graph of  $f$  is  $-1$  for  $x < 4$ ,  $1$  for  $x > 4$ , and undefined at  $x = 4$ .



43. The slope of the graph of  $f$  is negative for  $x < 0$  and positive for  $x > 0$ . The slope is undefined at  $x = 0$ .




44. The slope is positive for  $-2 < x < 0$  and negative for  $0 < x < 2$ . The slope is undefined at  $x = \pm 2$ , and 0 at  $x = 0$ .




45. Answers will vary.

Sample answer:  $y = -x$



46. Answers will vary.

Sample answer:  $y = x$



47.  $g(4) = 5$  because the tangent line passes through  $(4, 5)$ .

$$g'(4) = \frac{5 - 0}{4 - 7} = -\frac{5}{3}$$

48.  $h(-1) = 4$  because the tangent line passes through  $(-1, 4)$ .

$$h'(-1) = \frac{6 - 4}{3 - (-1)} = \frac{2}{4} = \frac{1}{2}$$

49.  $f(x) = 5 - 3x$  and  $c = 1$


50.  $f(x) = x^2$  and  $c = -2$

51.  $f(x) = -x^2$  and  $c = 6$

52.  $f(x) = 2\sqrt{x}$  and  $c = 9$


53.  $f(0) = 2$  and  $f'(x) = -3, -\infty < x < \infty$

$$f(x) = -3x + 2$$



54.  $f(0) = 4, f'(0) = 0; f'(x) < 0$  for  $x < 0, f'(x) > 0$  for  $x > 0$

Answers will vary. Sample answer:  $f(x) = x^2 + 4$



55. Let  $(x_0, y_0)$  be a point of tangency on the graph of  $f$ .

By the limit definition for the derivative,

$f'(x) = 4 - 2x$ . The slope of the line through  $(2, 5)$  and

$(x_0, y_0)$  equals the derivative of  $f$  at  $x_0$ :

$$\frac{5 - y_0}{2 - x_0} = 4 - 2x_0$$

$$5 - y_0 = (2 - x_0)(4 - 2x_0)$$

$$5 - (4x_0 - x_0^2) = 8 - 8x_0 + 2x_0^2$$

$$0 = x_0^2 - 4x_0 + 3$$

$$0 = (x_0 - 1)(x_0 - 3) \Rightarrow x_0 = 1, 3$$

Therefore, the points of tangency are  $(1, 3)$  and


$(3, 3)$ , and the corresponding slopes are 2 and  $-2$ . The

equations of the tangent lines are:

$$y - 5 = 2(x - 2) \quad y - 5 = -2(x - 2)$$

$$y = 2x + 1$$

$$y = -2x + 9$$



56. Let  $(x_0, y_0)$  be a point of tangency on the graph of  $f$ . By

the limit definition for the derivative,  $f'(x) = 2x$ . The

slope of the line through  $(1, -3)$  and  $(x_0, y_0)$  equals the

derivative of  $f$  at  $x_0$ :

$$\frac{-3 - y_0}{1 - x_0} = 2x_0$$

$$-3 - y_0 = (1 - x_0)2x_0$$

$$-3 - x_0^2 = 2x_0 - 2x_0^2$$

$$x_0^2 - 2x_0 - 3 = 0$$

$$(x_0 - 3)(x_0 + 1) = 0 \Rightarrow x_0 = 3, -1$$

Therefore, the points of tangency are  $(3, 9)$  and


$(-1, 1)$ , and the corresponding slopes are 6 and  $-2$ . The

equations of the tangent lines are:

$$y + 3 = 6(x - 1) \quad y + 3 = -2(x - 1)$$

$$y = 6x - 9$$

$$y = -2x - 1$$



57. (a)  $f(x) = x^2$

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^2 - x^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^2 + 2x(\Delta x) + (\Delta x)^2 - x^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\Delta x(2x + \Delta x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (2x + \Delta x) = 2x \end{aligned}$$


At  $x = -1$ ,  $f'(-1) = -2$  and the tangent line is

$$y - 1 = -2(x + 1) \quad \text{or} \quad y = -2x - 1.$$

At  $x = 0$ ,  $f'(0) = 0$  and the tangent line is  $y = 0$ .

At  $x = 1$ ,  $f'(1) = 2$  and the tangent line is

$$y = 2x - 1.$$



For this function, the slopes of the tangent lines are always distinct for different values of  $x$ .

$$\begin{aligned} \text{(b) } g'(x) &= \lim_{\Delta x \rightarrow 0} \frac{g(x + \Delta x) - g(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^3 - x^3}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^3 + 3x^2(\Delta x) + 3x(\Delta x)^2 + (\Delta x)^3 - x^3}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\Delta x(3x^2 + 3x(\Delta x) + (\Delta x)^2)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (3x^2 + 3x(\Delta x) + (\Delta x)^2) = 3x^2 \end{aligned}$$


At  $x = -1$ ,  $g'(-1) = 3$  and the tangent line is

$$y + 1 = 3(x + 1) \quad \text{or} \quad y = 3x + 2.$$

At  $x = 0$ ,  $g'(0) = 0$  and the tangent line is  $y = 0$ .

At  $x = 1$ ,  $g'(1) = 3$  and the tangent line is

$$y - 1 = 3(x - 1) \quad \text{or} \quad y = 3x - 2.$$



For this function, the slopes of the tangent lines are sometimes the same.

58. (a)  $g'(0) = -3$

(b)  $g'(3) = 0$


(c) Because  $g'(1) = -\frac{8}{3}$ ,  $g$  is decreasing (falling) at  $x = 1$ .

(d) Because  $g'(-4) = \frac{7}{3}$ ,  $g$  is increasing (rising) at  $x = -4$ .

(e) Because  $g'(4)$  and  $g'(6)$  are both positive,  $g(6)$  is greater than  $g(4)$ , and  $g(6) - g(4) > 0$ .

(f) No, it is not possible. All you can say is that  $g$  is decreasing (falling) at  $x = 2$ .


59.  $f(x) = \frac{1}{2}x^2$



$$f'(0) = 0, f'(1/2) = 1/2, f'(1) = 1, f'(2) = 2$$

(b) By symmetry:  $f'(-1/2) = -1/2, f'(-1) = -1, f'(-2) = -2$


(c)



$$(d) f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\frac{1}{2}(x + \Delta x)^2 - \frac{1}{2}x^2}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\frac{1}{2}(x^2 + 2x(\Delta x) + (\Delta x)^2) - \frac{1}{2}x^2}{\Delta x} = \lim_{\Delta x \rightarrow 0} \left( x + \frac{\Delta x}{2} \right) = x$$

60.  $f(x) = \frac{1}{3}x^3$


(a)



$$f'(0) = 0, f'(1/2) = 1/4, f'(1) = 1, f'(2) = 4, f'(3) = 9$$


 (b) By symmetry:  $f'(-1/2) = 1/4, f'(-1) = 1, f'(-2) = 4, f'(-3) = 9$ 

(c)




$$(d) f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ = \lim_{\Delta x \rightarrow 0} \frac{\frac{1}{3}(x + \Delta x)^3 - \frac{1}{3}x^3}{\Delta x} \\ = \lim_{\Delta x \rightarrow 0} \frac{\frac{1}{3}(x^3 + 3x^2(\Delta x) + 3x(\Delta x)^2 + (\Delta x)^3) - \frac{1}{3}x^3}{\Delta x} \\ = \lim_{\Delta x \rightarrow 0} \left[ x^2 + x(\Delta x) + \frac{1}{3}(\Delta x)^2 \right] = x^2$$

61.  $g(x) = \frac{f(x + 0.01) - f(x)}{0.01} \\ = \frac{[2(x + 0.01) - (x + 0.01)^2 - 2x + x^2]100}{0.01} \\ = 2 - 2x - 0.01$



The graph of  $g(x)$  is approximately the graph of  $f'(x) = 2 - 2x$ .

62.  $g(x) = \frac{f(x + 0.01) - f(x)}{0.01} \\ = \frac{(3\sqrt{x + 0.01} - 3\sqrt{x})100}{0.01}$



The graph of  $g(x)$  is approximately the graph of

$$f'(x) = \frac{3}{2\sqrt{x}}$$

63.  $f(2) = 2(4 - 2) = 4, f(2.1) = 2.1(4 - 2.1) = 3.99$

$$f'(2) \approx \frac{3.99 - 4}{2.1 - 2} = -0.1 \quad [\text{Exact: } f'(2) = 0]$$

64.  $f(2) = \frac{1}{4}(2^3) = 2, f(2.1) = 2.31525$

$$f'(2) \approx \frac{2.31525 - 2}{2.1 - 2} = 3.1525 \quad [\text{Exact: } f'(2) = 3]$$

65.  $f(x) = x^2 - 5, c = 3$

$$\begin{aligned} f'(3) &= \lim_{x \rightarrow 3} \frac{f(x) - f(3)}{x - 3} \\ &= \lim_{x \rightarrow 3} \frac{x^2 - 5 - (9 - 5)}{x - 3} \\ &= \lim_{x \rightarrow 3} \frac{(x - 3)(x + 3)}{x - 3} \\ &= \lim_{x \rightarrow 3} (x + 3) = 6 \end{aligned}$$

66.  $g(x) = x^2 - x, c = 1$

$$\begin{aligned} g'(1) &= \lim_{x \rightarrow 1} \frac{g(x) - g(1)}{x - 1} \\ &= \lim_{x \rightarrow 1} \frac{x^2 - x - 0}{x - 1} \\ &= \lim_{x \rightarrow 1} \frac{x(x - 1)}{x - 1} \\ &= \lim_{x \rightarrow 1} x = 1 \end{aligned}$$

67.  $f(x) = x^3 + 2x^2 + 1, c = -2$

$$\begin{aligned} f'(-2) &= \lim_{x \rightarrow -2} \frac{f(x) - f(-2)}{x - (-2)} \\ &= \lim_{x \rightarrow -2} \frac{(x^3 + 2x^2 + 1) - 1}{x + 2} \\ &= \lim_{x \rightarrow -2} \frac{x^2(x + 2)}{x + 2} = \lim_{x \rightarrow -2} x^2 = 4 \end{aligned}$$

68.  $f(x) = x^3 + 6x, c = 2$

$$\begin{aligned} f'(2) &= \lim_{x \rightarrow 2} \frac{f(x) - f(2)}{x - 2} \\ &= \lim_{x \rightarrow 2} \frac{(x^3 + 6x) - 20}{x - 2} \\ &= \lim_{x \rightarrow 2} \frac{(x - 2)(x^2 + 2x + 10)}{x - 2} \\ &= \lim_{x \rightarrow 2} (x^2 + 2x + 10) = 18 \end{aligned}$$

69.  $g(x) = \sqrt{|x|}, c = 0$

$$g'(0) = \lim_{x \rightarrow 0} \frac{g(x) - g(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{\sqrt{|x|}}{x}. \text{ Does not exist.}$$

$$\text{As } x \rightarrow 0^-, \frac{\sqrt{|x|}}{x} = \frac{-1}{\sqrt{|x|}} \rightarrow -\infty.$$

$$\text{As } x \rightarrow 0^+, \frac{\sqrt{|x|}}{x} = \frac{1}{\sqrt{x}} \rightarrow \infty.$$

Therefore  $g(x)$  is not differentiable at  $x = 0$ .

70.  $f(x) = \frac{3}{x}, c = 4$

$$\begin{aligned} f'(4) &= \lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4} \\ &= \lim_{x \rightarrow 4} \frac{\frac{3}{x} - \frac{3}{4}}{x - 4} \\ &= \lim_{x \rightarrow 4} \frac{12 - 3x}{4x(x - 4)} \\ &= \lim_{x \rightarrow 4} \frac{-3(x - 4)}{4x(x - 4)} \\ &= \lim_{x \rightarrow 4} \frac{3}{4x} = \frac{3}{16} \end{aligned}$$

71.  $f(x) = (x - 6)^{2/3}, c = 6$

$$\begin{aligned} f'(6) &= \lim_{x \rightarrow 6} \frac{f(x) - f(6)}{x - 6} \\ &= \lim_{x \rightarrow 6} \frac{(x - 6)^{2/3} - 0}{x - 6} = \lim_{x \rightarrow 6} \frac{1}{(x - 6)^{1/3}} \end{aligned}$$

Does not exist.

Therefore  $f(x)$  is not differentiable at  $x = 6$ .

72.  $g(x) = (x + 3)^{1/3}, c = -3$

$$\begin{aligned} g'(-3) &= \lim_{x \rightarrow -3} \frac{g(x) - g(-3)}{x - (-3)} \\ &= \lim_{x \rightarrow -3} \frac{(x + 3)^{1/3} - 0}{x + 3} = \lim_{x \rightarrow -3} \frac{1}{(x + 3)^{2/3}} \end{aligned}$$

Does not exist.

Therefore  $g(x)$  is not differentiable at  $x = -3$ .

73.  $h(x) = |x + 7|, c = -7$

$$\begin{aligned} h'(-7) &= \lim_{x \rightarrow -7} \frac{h(x) - h(-7)}{x - (-7)} \\ &= \lim_{x \rightarrow -7} \frac{|x + 7| - 0}{x + 7} = \lim_{x \rightarrow -7} \frac{|x + 7|}{x + 7} \end{aligned}$$

Does not exist.

Therefore  $h(x)$  is not differentiable at  $x = -7$ .

74.  $f(x) = |x - 6|, c = 6$

$$\begin{aligned} f'(6) &= \lim_{x \rightarrow 6} \frac{f(x) - f(6)}{x - 6} \\ &= \lim_{x \rightarrow 6} \frac{|x - 6| - 0}{x - 6} = \lim_{x \rightarrow 6} \frac{|x - 6|}{x - 6} \end{aligned}$$

Does not exist.

Therefore  $f(x)$  is not differentiable at  $x = 6$ .

75.  $f(x)$  is differentiable everywhere except at  $x = 3$ . (Discontinuity)

76.  $f(x)$  is differentiable everywhere except at  $x = \pm 3$ . (Sharp turns in the graph)


77.  $f(x)$  is differentiable everywhere except at  $x = -4$ . (Sharp turn in the graph)

78.  $f(x)$  is differentiable everywhere except at  $x = \pm 2$ . (Discontinuities)


79.  $f(x)$  is differentiable on the interval  $(1, \infty)$ . (At  $x = 1$  the tangent line is vertical.)

80.  $f(x)$  is differentiable everywhere except at  $x = 0$ . (Discontinuity)


81.  $f(x) = |x - 5|$  is differentiable everywhere except at  $x = -5$ . There is a sharp corner at  $x = 5$ .



82.  $f(x) = \frac{4x}{x - 3}$  is differentiable everywhere except at  $x = 3$ .  $f$  is not defined at  $x = 3$ . (Vertical asymptote)




83.  $f(x) = x^{2/5}$  is differentiable for all  $x \neq 0$ . There is a sharp corner at  $x = 0$ .



84.  $f$  is differentiable for all  $x \neq 1$ .

$f$  is not continuous at  $x = 1$ .



85.  $f(x) = |x - 1|$

The derivative from the left is

$$\lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1^-} \frac{|x - 1| - 0}{x - 1} = -1.$$

The derivative from the right is

$$\lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1^+} \frac{|x - 1| - 0}{x - 1} = 1.$$

The one-sided limits are not equal. Therefore,  $f$  is not differentiable at  $x = 1$ .

86.  $f(x) = \sqrt{1 - x^2}$

The derivative from the left does not exist because

$$\begin{aligned} \lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} &= \lim_{x \rightarrow 1^-} \frac{\sqrt{1 - x^2} - 0}{x - 1} \\ &= \lim_{x \rightarrow 1^-} \frac{\sqrt{1 - x^2}}{x - 1} \cdot \frac{\sqrt{1 - x^2}}{\sqrt{1 - x^2}} \\ &= \lim_{x \rightarrow 1^-} \frac{1 + x}{\sqrt{1 - x^2}} = -\infty. \end{aligned}$$

(Vertical tangent)

The limit from the right does not exist since  $f$  is undefined for  $x > 1$ . Therefore,  $f$  is not differentiable at  $x = 1$ .

87.  $f(x) = \begin{cases} (x - 1)^3, & x \leq 1 \\ (x - 1)^2, & x > 1 \end{cases}$

The derivative from the left is

$$\begin{aligned} \lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} &= \lim_{x \rightarrow 1^-} \frac{(x - 1)^3 - 0}{x - 1} \\ &= \lim_{x \rightarrow 1^-} (x - 1)^2 = 0. \end{aligned}$$

The derivative from the right is

$$\begin{aligned} \lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{x - 1} &= \lim_{x \rightarrow 1^+} \frac{(x - 1)^2 - 0}{x - 1} \\ &= \lim_{x \rightarrow 1^+} (x - 1) = 0. \end{aligned}$$

The one-sided limits are equal. Therefore,  $f$  is differentiable at  $x = 1$ . ( $f'(1) = 0$ )

$$88. f(x) = \begin{cases} x, & x \leq 1 \\ x^2, & x > 1 \end{cases}$$

The derivative from the left is

$$\lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1^-} \frac{x - 1}{x - 1} = \lim_{x \rightarrow 1^-} 1 = 1.$$

The derivative from the right is

$$\lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1^+} \frac{x^2 - 1}{x - 1} = \lim_{x \rightarrow 1^+} (x + 1) = 2.$$

The one-sided limits are not equal. Therefore,  $f$  is not differentiable at  $x = 1$ .

89. Note that  $f$  is continuous at  $x = 2$ .

$$f(x) = \begin{cases} x^2 + 1, & x \leq 2 \\ 4x - 3, & x > 2 \end{cases}$$

The derivative from the left is

$$\begin{aligned} \lim_{x \rightarrow 2^-} \frac{f(x) - f(2)}{x - 2} &= \lim_{x \rightarrow 2^-} \frac{(x^2 + 1) - 5}{x - 2} \\ &= \lim_{x \rightarrow 2^-} (x + 2) = 4. \end{aligned}$$

The derivative from the right is

$$\lim_{x \rightarrow 2^+} \frac{f(x) - f(2)}{x - 2} = \lim_{x \rightarrow 2^+} \frac{(4x - 3) - 5}{x - 2} = \lim_{x \rightarrow 2^+} 4 = 4.$$

The one-sided limits are equal. Therefore,  $f$  is differentiable at  $x = 2$ . ( $f'(2) = 4$ )

90. Note that  $f$  is continuous at  $x = 2$ .

$$f(x) = \begin{cases} \frac{1}{2}x + 1, & x < 2 \\ \sqrt{2x}, & x \geq 2 \end{cases}$$

The derivative from the left is


$$\begin{aligned} \lim_{x \rightarrow 2^-} \frac{f(x) - f(2)}{x - 2} &= \lim_{x \rightarrow 2^-} \frac{\left(\frac{1}{2}x + 1\right) - 2}{x - 2} \\ &= \lim_{x \rightarrow 2^-} \frac{\frac{1}{2}(x - 2)}{x - 2} = \frac{1}{2}. \end{aligned}$$

The derivative from the right is


$$\begin{aligned} \lim_{x \rightarrow 2^+} \frac{f(x) - f(2)}{x - 2} &= \lim_{x \rightarrow 2^+} \frac{\sqrt{2x} - 2}{x - 2} \cdot \frac{\sqrt{2x} + 2}{\sqrt{2x} + 2} \\ &= \lim_{x \rightarrow 2^+} \frac{2x - 4}{(x - 2)(\sqrt{2x} + 2)} \\ &= \lim_{x \rightarrow 2^+} \frac{2(x - 2)}{(x - 2)(\sqrt{2x} + 2)} \\ &= \lim_{x \rightarrow 2^+} \frac{2}{\sqrt{2x} + 2} = \frac{1}{2}. \end{aligned}$$

The one-sided limits are equal. Therefore,  $f$  is differentiable at  $x = 2$ . ( $f'(2) = \frac{1}{2}$ )

$$91. (a) \text{ The distance from } (3, 1) \text{ to the line } mx - y + 4 = 0 \text{ is } d = \frac{|Ax_1 + By_1 + C|}{\sqrt{A^2 + B^2}} = \frac{|m(3) - 1(1) + 4|}{\sqrt{m^2 + 1}} = \frac{|3m + 3|}{\sqrt{m^2 + 1}}$$




(b)




The function  $d$  is not differentiable at  $m = -1$ . This corresponds to the line  $y = -x + 4$ , which passes through the point  $(3, 1)$ .

92. (a)  $f(x) = x^2$  and  $f'(x) = 2x$



(b)  $g(x) = x^3$  and  $g'(x) = 3x^2$

(c) The derivative is a polynomial of degree 1 less than the original function. If  $h(x) = x^n$ , then  $h'(x) = nx^{n-1}$ .(d) If  $f(x) = x^4$ , then

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^4 - x^4}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^4 + 4x^3(\Delta x) + 6x^2(\Delta x)^2 + 4x(\Delta x)^3 + (\Delta x)^4 - x^4}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\Delta x(4x^3 + 6x^2(\Delta x) + 4x(\Delta x)^2 + (\Delta x)^3)}{\Delta x} = \lim_{\Delta x \rightarrow 0} (4x^3 + 6x^2(\Delta x) + 4x(\Delta x)^2 + (\Delta x)^3) = 4x^3. \end{aligned}$$

So, if  $f(x) = x^4$ , then  $f'(x) = 4x^3$  which is consistent with the conjecture. However, this is not a proof because you must verify the conjecture for all integer values of  $n$ ,  $n \geq 2$ .

93. False. The slope is  $\lim_{\Delta x \rightarrow 0} \frac{f(2 + \Delta x) - f(2)}{\Delta x}$ .

94. False.  $y = |x - 2|$  is continuous at  $x = 2$ , but is not differentiable at  $x = 2$ . (Sharp turn in the graph)95. False. If the derivative from the left of a point does not equal the derivative from the right of a point, then the derivative does not exist at that point. For example, if  $f(x) = |x|$ , then the derivative from the left at  $x = 0$  is  $-1$  and the derivative from the right at  $x = 0$  is  $1$ . At  $x = 0$ , the derivative does not exist.

96. True—see Theorem 2.1.



$$97. f(x) = \begin{cases} x \sin(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Using the Squeeze Theorem, you have  $-|x| \leq x \sin(1/x) \leq |x|$ ,  $x \neq 0$ . So,  $\lim_{x \rightarrow 0} x \sin(1/x) = 0 = f(0)$  and  $f$  is continuous at  $x = 0$ . Using the alternative form of the derivative, you have

$$\lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{x \sin(1/x) - 0}{x - 0} = \lim_{x \rightarrow 0} \left( \sin \frac{1}{x} \right).$$

Because this limit does not exist ( $\sin(1/x)$  oscillates between  $-1$  and  $1$ ), the function is not differentiable at  $x = 0$ .


$$g(x) = \begin{cases} x^2 \sin(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Using the Squeeze Theorem again, you have  $-x^2 \leq x^2 \sin(1/x) \leq x^2$ ,  $x \neq 0$ . So,  $\lim_{x \rightarrow 0} x^2 \sin(1/x) = 0 = g(0)$  and  $g$  is continuous at  $x = 0$ . Using the alternative form of the derivative again, you have

$$\lim_{x \rightarrow 0} \frac{g(x) - g(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{x^2 \sin(1/x) - 0}{x - 0} = \lim_{x \rightarrow 0} x \sin \frac{1}{x} = 0.$$

Therefore,  $g$  is differentiable at  $x = 0$ ,  $g'(0) = 0$ .

98.



As you zoom in, the graph of  $y_1 = x^2 + 1$  appears to be locally the graph of a horizontal line, whereas the graph of  $y_2 = |x| + 1$  always has a sharp corner at  $(0, 1)$ .  $y_2$  is not differentiable at  $(0, 1)$ .

## Section 2.2 Basic Differentiation Rules and Rates of Change

$$1. (a) \quad y = x^{1/2} \\ y' = \frac{1}{2}x^{-1/2} \\ y'(1) = \frac{1}{2}$$

$$(b) \quad y = x^3 \\ y' = 3x^2 \\ y'(1) = 3$$

$$2. (a) \quad y = x^{-1/2} \\ y' = -\frac{1}{2}x^{-3/2} \\ y'(1) = -\frac{1}{2}$$

$$(b) \quad y = x^{-1} \\ y' = -x^{-2} \\ y'(1) = -1$$

$$3. \quad y = 12 \\ y' = 0$$

$$4. \quad f(x) = -9 \\ f'(x) = 0$$

$$5. \quad y = x^7 \\ y' = 7x^6$$

$$6. \quad y = x^{12} \\ y' = 12x^{11}$$

$$7. \quad y = \frac{1}{x^5} = x^{-5} \\ y' = -5x^{-6} = -\frac{5}{x^6}$$

$$8. \quad y = \frac{3}{x^7} = 3x^{-7} \\ y' = 3(-7x^{-8}) = -\frac{21}{x^8}$$

