

# CHAPTER 7

## Applications of Integration

### Section 7.1 Area of a Region Between Two Curves

$$1. A = \int_0^6 [0 - (x^2 - 6x)] dx = -\int_0^6 (x^2 - 6x) dx$$

$$2. A = \int_{-2}^2 [(2x + 5) - (x^2 + 2x + 1)] dx$$

$$= \int_{-2}^2 (-x^2 + 4) dx$$

$$3. A = \int_0^3 [(-x^2 + 2x + 3) - (x^2 - 4x + 3)] dx$$

$$= \int_0^3 (-2x^2 + 6x) dx$$


$$4. A = \int_0^1 (x^2 - x^3) dx$$

$$5. A = 2 \int_{-1}^0 3(x^3 - x) dx = 6 \int_{-1}^0 (x^3 - x) dx$$


$$\text{or } -6 \int_0^1 (x^3 - x) dx$$

$$6. A = 2 \int_0^1 [(x - 1)^3 - (x - 1)] dx$$


$$7. \int_0^4 \left[ (x + 1) - \frac{x}{2} \right] dx$$




$$8. \int_{-1}^1 [(2 - x^2) - x^2] dx$$




$$9. \int_2^3 \left[ \left( \frac{x^3}{3} - x \right) - \frac{x}{3} \right] dx$$




$$10. \int_{-\pi/4}^{\pi/4} (\sec^2 x - \cos x) dx$$



$$11. \int_{-2}^1 [(2 - y) - y^2] dy$$



$$12. \int_0^4 (2\sqrt{y} - y) dy$$




23. The points of intersection are given by:

$$\sqrt{x} + 3 = \frac{1}{2}x + 3$$

$$\sqrt{x} = \frac{1}{2}x$$

$$x = \frac{x^2}{4} \quad \text{when } x = 0, 4$$



$$\begin{aligned} A &= \int_0^4 \left[ (\sqrt{x} + 3) - \left( \frac{1}{2}x + 3 \right) \right] dx \\ &= \left[ \frac{2}{3}x^{3/2} - \frac{x^2}{4} \right]_0^4 = \frac{16}{3} - 4 = \frac{4}{3} \end{aligned}$$

24. The points of intersection are given by:


$$\sqrt[3]{x-1} = x-1$$

$$x-1 = (x-1)^3 = x^3 - 3x^2 + 3x - 1$$

$$x^3 - 3x^2 + 2x = 0$$

$$x(x^2 - 3x + 2) = 0$$

$$x(x-2)(x-1) = 0 \quad \text{when } x = 0, 1, 2$$




$$\begin{aligned} A &= 2 \int_0^1 \left[ (x-1) - \sqrt[3]{x-1} \right] dx \\ &= 2 \left[ \frac{x^2}{2} - x - \frac{3}{4}(x-1)^{4/3} \right]_0^1 \\ &= 2 \left[ \left( \frac{1}{2} - 1 - 0 \right) - \left( -\frac{3}{4} \right) \right] = \frac{1}{2} \end{aligned}$$

25. The points of intersection are given by:

$$y^2 = y + 2$$

$$(y-2)(y+1) = 0 \quad \text{when } y = -1, 2$$




$$\begin{aligned} A &= \int_{-1}^2 [g(y) - f(y)] dy \\ &= \int_{-1}^2 [(y+2) - y^2] dy \\ &= \left[ 2y + \frac{y^2}{2} - \frac{y^3}{3} \right]_{-1}^2 = \frac{9}{2} \end{aligned}$$

26. The points of intersection are given by:


$$2y - y^2 = -y$$

$$y(y-3) = 0 \quad \text{when } y = 0, 3$$




$$\begin{aligned} A &= \int_0^3 [f(y) - g(y)] dy \\ &= \int_0^3 [(2y - y^2) - (-y)] dy \\ &= \int_0^3 (3y - y^2) dy \\ &= \left[ \frac{3}{2}y^2 - \frac{1}{3}y^3 \right]_0^3 = \frac{9}{2} \end{aligned}$$

27.




$$\begin{aligned} A &= \int_{-1}^2 [f(y) - g(y)] dy \\ &= \int_{-1}^2 [(y^2 + 1) - 0] dy \\ &= \left[ \frac{y^3}{3} + y \right]_{-1}^2 = 6 \end{aligned}$$

28.



$$\begin{aligned}
 A &= \int_0^3 [f(y) - g(y)] dy \\
 &= \int_0^3 \left[ \frac{y}{\sqrt{16 - y^2}} - 0 \right] dy \\
 &= -\frac{1}{2} \int_0^3 (16 - y^2)^{-1/2} (-2y) dy \\
 &= \left[ -\sqrt{16 - y^2} \right]_0^3 = 4 - \sqrt{7} \approx 1.354
 \end{aligned}$$


29.  $y = \frac{10}{x} \Rightarrow x = \frac{10}{y}$



$$\begin{aligned}
 A &= \int_2^{10} \frac{10}{y} dy \\
 &= [10 \ln y]_2^{10} \\
 &= 10(\ln 10 - \ln 2) \\
 &= 10 \ln 5 \approx 16.0944
 \end{aligned}$$


30. The point of intersection is given by:

$$\begin{aligned}
 \frac{4}{2 - x} &= 4 \\
 \frac{4}{2 - x} - 4 &= 0 \quad \text{when } x = 1
 \end{aligned}$$



$$\begin{aligned}
 A &= \int_0^1 \left( 4 - \frac{4}{2 - x} \right) dx \\
 &= [4x + 4 \ln |2 - x|]_0^1 \\
 &= 4 - 4 \ln 2 \\
 &\approx 1.227
 \end{aligned}$$

31. (a)




(b) The points of intersection are given by:

$$\begin{aligned}
 x^3 - 3x^2 + 3x &= x^2 \\
 x(x - 1)(x - 3) &= 0 \quad \text{when } x = 0, 1, 3
 \end{aligned}$$

$$\begin{aligned}
 A &= \int_0^1 [f(x) - g(x)] dx + \int_1^3 [g(x) - f(x)] dx \\
 &= \int_0^1 [(x^3 - 3x^2 + 3x) - x^2] dx + \int_1^3 [x^2 - (x^3 - 3x^2 + 3x)] dx \\
 &= \int_0^1 (x^3 - 4x^2 + 3x) dx + \int_1^3 (-x^3 + 4x^2 - 3x) dx = \left[ \frac{x^4}{4} - \frac{4}{3}x^3 + \frac{3}{2}x^2 \right]_0^1 + \left[ -\frac{x^4}{4} + \frac{4}{3}x^3 - \frac{3}{2}x^2 \right]_1^3 = \frac{5}{12} + \frac{8}{3} = \frac{37}{12}
 \end{aligned}$$

 (c) Numerical approximation:  $0.417 + 2.667 \approx 3.083$ 

32. (a)




(b) The points of intersection are given by:

$$\begin{aligned}
 x^4 - 2x^2 &= 2x^2 \\
 x^2(x^2 - 4) &= 0 \quad \text{when } x = 0, \pm 2
 \end{aligned}$$

$$A = 2 \int_0^2 [2x^2 - (x^4 - 2x^2)] dx = 2 \int_0^2 (4x^2 - x^4) dx = 2 \left[ \frac{4x^3}{3} - \frac{x^5}{5} \right]_0^2 = \frac{128}{15}$$

(c) Numerical approximation: 8.533

33. (a)  $f(x) = x^4 - 4x^2$ ,  $g(x) = x^2 - 4$



(b) The points of intersection are given by:

$$x^4 - 4x^2 = x^2 - 4$$

$$x^4 - 5x^2 + 4 = 0$$

$$(x^2 - 4)(x^2 - 1) = 0 \quad \text{when } x = \pm 2, \pm 1$$


By symmetry:

$$\begin{aligned} A &= 2 \int_0^1 [(x^4 - 4x^2) - (x^2 - 4)] dx + 2 \int_1^2 [(x^2 - 4) - (x^4 - 4x^2)] dx \\ &= 2 \int_0^1 (x^4 - 5x^2 + 4) dx + 2 \int_1^2 (-x^4 + 5x^2 - 4) dx \\ &= 2 \left[ \frac{x^5}{5} - \frac{5x^3}{3} + 4x \right]_0^1 + 2 \left[ -\frac{x^5}{5} + \frac{5x^3}{3} - 4x \right]_1^2 \\ &= 2 \left[ \frac{1}{5} - \frac{5}{3} + 4 \right] + 2 \left[ \left( -\frac{32}{5} + \frac{40}{3} - 8 \right) - \left( -\frac{1}{5} + \frac{5}{3} - 4 \right) \right] = 8 \end{aligned}$$

(c) Numerical approximation:

$$5.067 + 2.933 = 8.0$$

34. (a)



(b) The points of intersection are given by:

$$x^4 - 9x^2 = x^3 - 9x$$


$$x^4 - x^3 - 9x^2 + 9x = 0$$

$$x(x - 3)(x - 1)(x + 3) = 0 \quad \text{when } x = -3, 0, 1, 3$$

$$\begin{aligned} A &= \int_{-3}^0 [(x^3 - 9x) - (x^4 - 9x^2)] dx + \int_0^1 [(x^4 - 9x^2) - (x^3 - 9x)] dx + \int_1^3 [(x^3 - 9x) - (x^4 - 9x^2)] dx \\ &= \left[ \frac{x^4}{4} - \frac{9x^2}{2} - \frac{x^5}{5} + 3x^3 \right]_{-3}^0 + \left[ \frac{x^5}{5} - 3x^3 - \frac{x^4}{4} + \frac{9x^2}{2} \right]_0^1 + \left[ \frac{x^4}{4} - \frac{9x^2}{2} - \frac{x^5}{5} + 3x^3 \right]_1^3 \\ &= \frac{1053}{20} + \frac{29}{20} + \frac{68}{5} = \frac{677}{10} \end{aligned}$$

(c) Numerical approximation: 67.7

35. (a)




(b) The points of intersection are given by:

$$\begin{aligned} \frac{1}{1+x^2} &= \frac{x^2}{2} \\ x^4 + x^2 - 2 &= 0 \\ (x^2 + 2)(x^2 - 1) &= 0 \text{ when } x = \pm 1 \\ A &= 2 \int_0^1 [f(x) - g(x)] dx \\ &= 2 \int_0^1 \left[ \frac{1}{1+x^2} - \frac{x^2}{2} \right] dx \\ &= 2 \left[ \arctan x - \frac{x^3}{6} \right]_0^1 \\ &= 2 \left( \frac{\pi}{4} - \frac{1}{6} \right) = \frac{\pi}{2} - \frac{1}{3} \approx 1.237 \end{aligned}$$

(c) Numerical approximation: 1.237


36. (a)




$$\begin{aligned} (b) \quad A &= \int_0^3 \left[ \frac{6x}{x^2 + 1} - 0 \right] dx \\ &= \left[ 3 \ln(x^2 + 1) \right]_0^3 \\ &= 3 \ln 10 \\ &\approx 6.908 \end{aligned}$$

(c) Numerical approximation: 6.908


$$\begin{aligned} 37. \quad A &= \int_0^{2\pi} [(2 - \cos x) - \cos x] dx \\ &= 2 \int_0^{2\pi} (1 - \cos x) dx \\ &= 2[x - \sin x]_0^{2\pi} = 4\pi \approx 12.566 \end{aligned}$$




$$\begin{aligned} 38. \quad A &= \int_{-\pi/2}^{\pi/6} (\cos 2x - \sin x) dx \\ &= \left[ \frac{1}{2} \sin 2x + \cos x \right]_{-\pi/2}^{\pi/6} \\ &= \left( \frac{\sqrt{3}}{4} + \frac{\sqrt{3}}{2} \right) - (0) = \frac{3\sqrt{3}}{4} \approx 1.299 \end{aligned}$$



$$\begin{aligned} 39. \quad A &= 2 \int_0^{\pi/3} [f(x) - g(x)] dx \\ &= 2 \int_0^{\pi/3} (2 \sin x - \tan x) dx \\ &= 2[-2 \cos x + \ln|\cos x|]_0^{\pi/3} = 2(1 - \ln 2) \approx 0.614 \end{aligned}$$




$$\begin{aligned} 40. \quad A &= \int_0^1 \left[ (\sqrt{2} - 4)x + 4 - \sec \frac{\pi x}{4} \tan \frac{\pi x}{4} \right] dx \\ &= \left[ \frac{\sqrt{2} - 4}{2} x^2 + 4x - \frac{4}{\pi} \sec \frac{\pi x}{4} \right]_0^1 \\ &= \left( \frac{\sqrt{2} - 4}{2} + 4 - \frac{4}{\pi} \sqrt{2} \right) - \left( -\frac{4}{\pi} \right) \\ &= \frac{\sqrt{2}}{2} + 2 + \frac{4}{\pi} (1 - \sqrt{2}) \approx 2.1797 \end{aligned}$$



$$41. A = \int_0^1 [xe^{-x^2} - 0] dx$$

$$= \left[ -\frac{1}{2}e^{-x^2} \right]_0^1 = \frac{1}{2} \left( 1 - \frac{1}{e} \right) \approx 0.316$$





42. From the graph,  $f$  and  $g$  intersect at  $x = 0$  and  $x = 2$ .

$$A = \int_0^2 \left[ \left( \frac{3}{2}x + 1 \right) - 2^x \right] dx$$

$$= \left[ \frac{3x^2}{4} + x - \frac{2^x}{\ln 2} \right]_0^2$$

$$= \left( 3 + 2 - \frac{4}{\ln 2} \right) + \frac{1}{\ln 2}$$

$$= 5 - \frac{3}{\ln 2} \approx 0.672$$




(b)  $A = \int_0^\pi (2 \sin x + \sin 2x) dx$

$$= \left[ -2 \cos x - \frac{1}{2} \cos 2x \right]_0^\pi$$

$$= \left( 2 - \frac{1}{2} \right) - \left( -2 - \frac{1}{2} \right) = 4$$


(c) Numerical approximation: 4.0



(b)  $A = \int_0^\pi (2 \sin x + \cos 2x) dx$

$$= \left[ -2 \cos x + \frac{1}{2} \sin 2x \right]_0^\pi = 4$$

(c) Numerical approximation: 4




(b)  $A = \int_1^3 \frac{1}{x^2} e^{1/x} dx$

$$= \left[ -e^{-1/x} \right]_1^3$$

$$= e - e^{1/3}$$

(c) Numerical approximation: 1.323




(b)  $A = \int_1^5 \frac{4 \ln x}{x} dx$

$$= \left[ 2(\ln x)^2 \right]_1^5$$

$$= 2(\ln 5)^2$$

(c) Numerical approximation: 5.181




(b) The integral

$$A = \int_0^3 \sqrt{\frac{x^3}{4-x}} dx$$

does not have an elementary antiderivative.

(c)  $A \approx 4.7721$




(b) The integral

$$A = \int_0^1 \sqrt{x} e^x dx$$

does not have an elementary antiderivative.

(c) 1.2556


49. (a)



(b) The intersection points are difficult to determine by hand.

(c)  $\text{Area} = \int_{-c}^c [4 \cos x - x^2] dx \approx 6.3043$  where  $c \approx 1.201538$ .

50. (a)




(b) The intersection points are difficult to determine.


 (c) Intersection points:  $(-1.164035, 1.3549778)$  and  $(1.4526269, 2.1101248)$ 

$$A = \int_{-1.164035}^{1.4526269} [\sqrt{3+x} - x^2] dx \approx 3.0578$$


51.  $F(x) = \int_0^x \left(\frac{1}{2}t + 1\right) dt = \left[\frac{t^2}{4} + t\right]_0^x = \frac{x^2}{4} + x$

 (a)  $F(0) = 0$ 



(b)  $F(2) = \frac{2^2}{4} + 2 = 3$




(c)  $F(6) = \frac{6^2}{4} + 6 = 15$




52.  $F(x) = \int_0^x \left(\frac{1}{2}t^2 + 2\right) dt = \left[\frac{1}{6}t^3 + 2t\right]_0^x = \frac{x^3}{6} + 2x$

 (a)  $F(0) = 0$ 


(b)  $F(4) = \frac{4^3}{6} + 2(4) = \frac{56}{3}$




(c)  $F(6) = 36 + 12 = 48$




$$53. F(\alpha) = \int_{-1}^{\alpha} \cos \frac{\pi\theta}{2} d\theta = \left[ \frac{2}{\pi} \sin \frac{\pi\theta}{2} \right]_{-1}^{\alpha} = \frac{2}{\pi} \sin \frac{\pi\alpha}{2} + \frac{2}{\pi}$$


(a)  $F(-1) = 0$



(b)  $F(0) = \frac{2}{\pi} \approx 0.6366$




(c)  $F\left(\frac{1}{2}\right) = \frac{2 + \sqrt{2}}{\pi} \approx 1.0868$




$$54. F(y) = \int_{-1}^y 4e^{x/2} dx = \left[ 8e^{x/2} \right]_{-1}^y = 8e^{y/2} - 8e^{-1/2}$$


(a)  $F(-1) = 0$



(b)  $F(0) = 8 - 8e^{-1/2} \approx 3.1478$




(c)  $F(4) = 8e^2 - 8e^{-1/2} \approx 54.2602$



$$55. A = \int_2^4 \left[ \left( \frac{3}{2}x - 12 \right) - (x - 5) \right] dx + \int_4^6 \left[ \left( -\frac{5}{2}x + 16 \right) - (x - 5) \right] dx$$

$$= \int_2^4 \left( \frac{1}{2}x - 7 \right) dx + \int_4^6 \left( -\frac{7}{2}x + 21 \right) dx = \left[ \frac{1}{4}x^2 - 7x \right]_2^4 + \left[ -\frac{7}{4}x^2 + 21x \right]_4^6 = 7 + 7 = 14$$




$$56. A = \int_0^4 \frac{3}{4}x dx + \int_4^6 \left( 9 - \frac{3}{2}x \right) dx$$

$$= \left[ \frac{3x^2}{8} \right]_0^4 + \left[ 9x - \frac{3x^2}{4} \right]_4^6$$


$$= 6 + (54 - 27) - (36 - 12)$$

$$= 6 + 3 = 9$$





57.


 Left boundary line:  $y = x + 2 \Leftrightarrow x = y - 2$ 

 Right boundary line:  $y = x - 2 \Leftrightarrow x = y + 2$ 

$$\begin{aligned} A &= \int_{-2}^2 [(y + 2) - (y - 2)] dy \\ &= \int_{-2}^2 4 dy = [4y]_{-2}^2 = 8 - (-8) = 16 \end{aligned}$$


$$58. A = \int_0^1 [2x - (-3x)] dx + \int_1^3 [(-2x + 4) - (\frac{1}{2}x - \frac{7}{2})] dx$$

$$= \int_0^1 5x dx + \int_1^3 \left(-\frac{5}{2}x + \frac{15}{2}\right) dx$$

$$= \left[\frac{5x^2}{2}\right]_0^1 + \left[-\frac{5x^2}{4} + \frac{15}{2}x\right]_1^3$$

$$= \frac{5}{2} + \left(-\frac{45}{4} + \frac{45}{2} + \frac{5}{4} - \frac{15}{2}\right)$$

$$= \frac{15}{2}$$


 59. Answers will vary. *Sample answer:* If you let  $\Delta x = 6$  and  $n = 10$ ,  $b - a = 10(6) = 60$ .

$$(a) \text{ Area} \approx \frac{60}{2(10)} [0 + 2(14) + 2(14) + 2(12) + 2(12) + 2(15) + 2(20) + 2(23) + 2(25) + 2(26) + 0] = 3[322] = 966 \text{ ft}^2$$

$$(b) \text{ Area} \approx \frac{60}{3(10)} [0 + 4(14) + 2(14) + 4(12) + 2(12) + 4(15) + 2(20) + 4(23) + 2(25) + 4(26) + 0] = 2[502] = 1004 \text{ ft}^2$$

 60. Answers will vary. *Sample answer:*  $\Delta x = 4$ ,  $n = 8$ ,  $b - a = (8)(4) = 32$ 

$$(a) \text{ Area} \approx \frac{32}{2(8)} [0 + 2(11) + 2(13.5) + 2(14.2) + 2(14) + 2(14.2) + 2(15) + 2(13.5) + 0]$$

$$= 2[190.8]$$

$$= 381.6 \text{ mi}^2$$

$$(b) \text{ Area} \approx \frac{32}{3(8)} [0 + 4(11) + 2(13.5) + 4(14.2) + 2(14) + 4(14.2) + 2(15) + 4(13.5) + 0]$$

$$= \frac{4}{3}[296.6]$$

$$= 395.5 \text{ mi}^2$$

61.  $f(x) = x^3$


$f'(x) = 3x^2$

 At  $(1, 1)$ ,  $f'(1) = 3$ .

 Tangent line:  $y - 1 = 3(x - 1)$  or  $y = 3x - 2$ 

 The tangent line intersects  $f(x) = x^3$  at  $x = -2$ .

$$A = \int_{-2}^1 [x^3 - (3x - 2)] dx = \left[\frac{x^4}{4} - \frac{3x^2}{2} + 2x\right]_{-2}^1 = \frac{27}{4}$$



62.  $y = x^3 - 2x, \quad (-1, 1)$

$$y' = 3x^2 - 2$$


$$y'(-1) = 3 - 2 = 1$$

$$\text{Tangent line: } y - 1 = 1(x + 1) \Rightarrow y = x + 2$$

Intersection points:  $(-1, 1)$  and  $(2, 4)$ 

$$A = \int_{-1}^2 [(x+2) - (x^3 - 2x)] dx = \int_{-1}^2 (-x^3 + 3x + 2) dx$$

$$= \left[ -\frac{x^4}{4} + \frac{3x^2}{2} + 2x \right]_{-1}^2 = \left[ (-4 + 6 + 4) - \left( -\frac{1}{4} + \frac{3}{2} - 2 \right) \right] = \frac{27}{4}$$



63.  $f(x) = \frac{1}{x^2 + 1}$


$$f'(x) = -\frac{2x}{(x^2 + 1)^2}$$

At  $\left(1, \frac{1}{2}\right)$ ,  $f'(1) = -\frac{1}{2}$

$$\text{Tangent line: } y - \frac{1}{2} = -\frac{1}{2}(x - 1) \text{ or } y = -\frac{1}{2}x + 1$$

The tangent line intersects  $f(x) = \frac{1}{x^2 + 1}$  at  $x = 0$ .

$$A = \int_0^1 \left[ \frac{1}{x^2 + 1} - \left( -\frac{1}{2}x + 1 \right) \right] dx = \left[ \arctan x + \frac{x^2}{4} - x \right]_0^1 = \frac{\pi - 3}{4} \approx 0.0354$$




64.  $y = \frac{2}{1 + 4x^2}, \quad \left(\frac{1}{2}, 1\right)$

$$y' = \frac{-16x}{(1 + 4x^2)^2}$$

$$y'\left(\frac{1}{2}\right) = \frac{-8}{2^2} = -2$$

$$\text{Tangent line: } y - 1 = -2\left(x - \frac{1}{2}\right)$$


$$y = -2x + 2$$

Intersection points:  $\left(\frac{1}{2}, 1\right), (0, 2)$ 

$$A = \int_0^{1/2} \left[ \frac{2}{1 + 4x^2} - (-2x + 2) \right] dx = \left[ \arctan(2x) + x^2 - 2x \right]_0^{1/2} = \arctan(1) + \frac{1}{4} - 1 = \frac{\pi}{4} - \frac{3}{4} \approx 0.0354$$

65.  $x^4 - 2x^2 + 1 \leq 1 - x^2$  on  $[-1, 1]$

$$\begin{aligned} A &= \int_{-1}^1 [(1 - x^2) - (x^4 - 2x^2 + 1)] dx \\ &= \int_{-1}^1 (x^2 - x^4) dx \\ &= \left[ \frac{x^3}{3} - \frac{x^5}{5} \right]_{-1}^1 = \frac{4}{15} \end{aligned}$$



You can use a single integral because  $x^4 - 2x^2 + 1 \leq 1 - x^2$  on  $[-1, 1]$ .


66.  $x^3 \geq x$  on  $[-1, 0]$ ,  $x^3 \leq x$  on  $[0, 1]$

Both functions symmetric to origin.

$$\int_{-1}^0 (x^3 - x) dx = -\int_0^1 (x^3 - x) dx$$

$$\text{Thus, } \int_{-1}^1 (x^3 - x) dx = 0.$$

$$A = 2 \int_0^1 (x - x^3) dx = 2 \left[ \frac{x^2}{2} - \frac{x^4}{4} \right]_0^1 = \frac{1}{2}$$



67. (a)  $\int_0^5 [v_1(t) - v_2(t)] dt = 10$  means that Car 1 traveled

10 more meters than Car 2 on the interval  $0 \leq t \leq 5$ .

$$\int_0^{10} [v_1(t) - v_2(t)] dt = 30 \text{ means that Car 1}$$

traveled 30 more meters than Car 2 on the interval  $0 \leq t \leq 10$ .

$$\int_{20}^{30} [v_1(t) - v_2(t)] dt = -5 \text{ means that Car 2}$$

traveled 5 more meters than Car 1 on the interval  $20 \leq t \leq 30$ .

(b) No, it is not possible because you do not know the initial distance between the cars.

(c) At  $t = 10$ , Car 1 is ahead by 30 meters.

(d) At  $t = 20$ , Car 1 is ahead of Car 2 by 13 meters.

From part (a), at  $t = 30$ , Car 1 is ahead by  $13 - 5 = 8$  meters.

68. (a) The area between the two curves represents the difference between the accumulated deficit under the two plans.

(b) Proposal 2 is better because the cumulative deficit (the area under the curve) is less.

69.  $A = \int_{-3}^3 (9 - x^2) dx = 36$

$$\int_{-\sqrt{9-b}}^{\sqrt{9-b}} [(9 - x^2) - b] dx = 18$$

$$\int_0^{\sqrt{9-b}} [(9 - b) - x^2] dx = 9$$


$$\left[ (9 - b)x - \frac{x^3}{3} \right]_0^{\sqrt{9-b}} = 9$$

$$\frac{2}{3}(9 - b)^{3/2} = 9$$


$$(9 - b)^{3/2} = \frac{27}{2}$$

$$9 - b = \frac{9}{\sqrt[3]{4}}$$

$$b = 9 - \frac{9}{\sqrt[3]{4}} \approx 3.330$$




$$\begin{aligned}
 70. \quad A &= 2 \int_0^9 (9-x) dx = 2 \left[ 9x - \frac{x^2}{2} \right]_0^9 = 81 \\
 2 \int_0^{9-b} [(9-x) - b] dx &= \frac{81}{2} \\
 2 \int_0^{9-b} [(9-b) - x] dx &= \frac{81}{2} \\
 2 \left[ (9-b)x - \frac{x^2}{2} \right]_0^{9-b} &= \frac{81}{2} \\
 (9-b)(9-b) &= \frac{81}{2} \\
 9-b &= \frac{9}{\sqrt{2}} \\
 b &= 9 - \frac{9}{\sqrt{2}} \approx 2.636
 \end{aligned}$$




71. Area of triangle  $OAB$  is  $\frac{1}{2}(4)(4) = 8$ .

$$\begin{aligned}
 4 &= \int_0^a (4-x) dx = \left[ 4x - \frac{x^2}{2} \right]_0^a = 4a - \frac{a^2}{2} \\
 a^2 - 8a + 8 &= 0 \\
 a &= 4 \pm 2\sqrt{2}
 \end{aligned}$$

Because  $0 < a < 4$ , select  $a = 4 - 2\sqrt{2} \approx 1.172$ .




$$\begin{aligned}
 72. \quad \text{Total area} &= \int_{-2}^2 (4-y^2) dy = 2 \int_0^2 (4-y^2) dy \\
 &= 2 \left[ 4y - \frac{y^3}{3} \right]_0^2 = 2 \left[ 8 - \frac{8}{3} \right] = \frac{32}{3} \\
 \frac{16}{3} &= 2 \int_a^4 \sqrt{4-x} dx = -\frac{4}{3} (4-x)^{3/2} \Big|_a^4 = \frac{4}{3} (4-a)^{3/2} \\
 4 &= (4-a)^{3/2} \\
 4^{2/3} &= 4-a \\
 a &= 4 - 4^{2/3} \approx 1.48
 \end{aligned}$$



$$73. \quad \lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n (x_i - x_i^2) \Delta x$$

where  $x_i = \frac{i}{n}$  and  $\Delta x = \frac{1}{n}$  is the same as


$$\int_0^1 (x - x^2) dx = \left[ \frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = \frac{1}{6}$$



$$74. \quad \lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n (4 - x_i^2) \Delta x$$

where  $x_i = -2 + \frac{4i}{n}$  and  $\Delta x = \frac{4}{n}$  is the same as

$$\int_{-2}^2 (4 - x^2) dx = \left[ 4x - \frac{x^3}{3} \right]_{-2}^2 = \frac{32}{3}$$





75.  $R_1$  projects the greater revenue because the area under the curve is greater.

$$\begin{aligned}
 &\int_{15}^{20} [(7.21 + 0.58t) - (7.21 + 0.45t)] dt \\
 &= \int_{15}^{20} 0.13t dt = \left[ \frac{0.13t^2}{2} \right]_{15}^{20} = \$11.375 \text{ billion}
 \end{aligned}$$

- 76.
- $R_2$
- projects the greater revenue because the area under the curve is greater.

$$\begin{aligned} & \int_{15}^{20} [(7.21 + 0.26t + 0.02t^2) - (7.21 + 0.1t + 0.01t^2)] dt \\ &= \int_{15}^{20} (0.01t^2 + 0.16t) dt \\ &= \left[ \frac{0.01t^3}{3} + \frac{0.16t^2}{2} \right]_{15}^{20} \approx \$29.417 \text{ billion} \end{aligned}$$

77. (a)
- $y_1 = 0.0124x^2 - 0.385x + 7.85$



(d) Income inequality =  $\int_0^{100} [x - y_1] dx \approx 2006.7$

78. 5%:
- $P_1 = 15.9e^{0.05t}$
- (in millions)

3.5%:  $P_2 = 15.9e^{0.035t}$  (in millions)

Difference in profits over 5 years:

$$\int_0^5 (P_1 - P_2) dt = \int_0^5 15.9(e^{0.05t} - e^{0.035t}) dt = 15.9 \left[ \frac{e^{0.05t}}{0.05} - \frac{e^{0.035t}}{0.035} \right]_0^5 \approx \$3.44 \text{ million}$$

79. (a)  $A = 2 \left[ \int_0^5 \left( 1 - \frac{1}{3}\sqrt{5-x} \right) dx + \int_5^{5.5} (1-0) dx \right]$

$$= 2 \left[ \left[ x + \frac{2}{9}(5-x)^{3/2} \right]_0^5 + [x]_5^{5.5} \right]$$

$$= 2 \left( 5 - \frac{10\sqrt{5}}{9} + 5.5 - 5 \right) \approx 6.031 \text{ m}^2$$

(b)  $V = 2A \approx 2(6.031) \approx 12.062 \text{ m}^3$

(c)  $5000V \approx 5000(12.062) = 60,310 \text{ pounds}$

80. The curves intersect at the point where the slope of
- $y_2$
- equals that of
- $y_1$
- , 1.

$$y_2 = 0.08x^2 + k \Rightarrow y_2' = 0.16x = 1 \Rightarrow x = \frac{1}{0.16} = 6.25$$

- (a) The value of
- $k$
- is given by

$$y_1 = y_2$$

$$6.25 = (0.08)(6.25)^2 + k$$

$$k = 3.125.$$

(b) Area =  $2 \int_0^{6.25} (y_2 - y_1) dx$

$$= 2 \int_0^{6.25} (0.08x^2 + 3.125 - x) dx$$

$$= 2 \left[ \frac{0.08x^3}{3} + 3.125x - \frac{x^2}{2} \right]_0^{6.25}$$

$$= 2(6.510417) \approx 13.02083$$


81. Line:  $y = \frac{-3}{7\pi}x$

$$A = \int_0^{7\pi/6} \left[ \sin x + \frac{3x}{7\pi} \right] dx$$

$$= \left[ -\cos x + \frac{3x^2}{14\pi} \right]_0^{7\pi/6}$$

$$= \frac{\sqrt{3}}{2} + \frac{7\pi}{24} + 1$$


$$\approx 2.7823$$



82.  $A = 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx = \frac{4b}{a} \int_0^a \sqrt{a^2 - x^2} dx$

$$\int_0^a \sqrt{a^2 - x^2} dx \text{ is the area of } \frac{1}{4} \text{ of a circle} = \frac{\pi a^2}{4}.$$

$$\text{So, } A = \frac{4b}{a} \left( \frac{\pi a^2}{4} \right) = \pi ab.$$

83. True. The region has been shifted  $C$  units upward (if  $C > 0$ ), or  $C$  units downward (if  $C < 0$ ).

84. True. This is a property of integrals.

85. False. Let  $f(x) = x$  and  $g(x) = 2x - x^2$ ,  $f$  and  $g$  intersect at  $(1, 1)$ , the midpoint of  $[0, 2]$ , but

$$\int_0^2 [f(x) - g(x)] dx = \int_0^2 [x - (2x - x^2)] dx = \frac{2}{3} \neq 0.$$

86. True. The area under  $f(x)$  between 0 and 1 is  $\frac{1}{6}$ . Thecurves intersect at  $x = \frac{1}{2}^{1/3}$ , and the area between  $y = \left(1 - \frac{1}{2}^{1/3}\right)x$  and  $f$  on the interval  $\left[0, \frac{1}{2}^{1/3}\right]$  is  $\frac{1}{12}$ .87. You want to find  $c$  such that:

$$\int_0^b [(2x - 3x^3) - c] dx = 0$$

$$\left[ x^2 - \frac{3}{4}x^4 - cx \right]_0^b = 0$$

$$b^2 - \frac{3}{4}b^4 - cb = 0$$

But,  $c = 2b - 3b^3$  because  $(b, c)$  is on the graph.


$$b^2 - \frac{3}{4}b^4 - (2b - 3b^3)b = 0$$

$$4 - 3b^2 - 8 + 12b^2 = 0$$

$$9b^2 = 4$$

$$b = \frac{2}{3}$$

$$c = \frac{4}{9}$$



## Section 7.2 Volume: The Disk Method

1.  $V = \pi \int_0^1 (-x + 1)^2 dx = \pi \int_0^1 (x^2 - 2x + 1) dx = \pi \left[ \frac{x^3}{3} - x^2 + x \right]_0^1 = \frac{\pi}{3}$

2.  $V = \pi \int_0^2 (4 - x^2)^2 dx = \pi \int_0^2 (x^4 - 8x^2 + 16) dx = \pi \left[ \frac{x^5}{5} - \frac{8x^3}{3} + 16x \right]_0^2 = \frac{256\pi}{15}$

3.  $V = \pi \int_1^4 (\sqrt{x})^2 dx = \pi \int_1^4 x dx = \pi \left[ \frac{x^2}{2} \right]_1^4 = \frac{15\pi}{2}$

4.  $V = \pi \int_0^3 (\sqrt{9 - x^2})^2 dx = \pi \int_0^3 (9 - x^2) dx = \pi \left[ 9x - \frac{x^3}{3} \right]_0^3 = 18\pi$