

7.4 Exercises


See CalcChat.com for tutorial help and worked-out solutions to odd-numbered exercises.

Finding Distance Using Two Methods In Exercises 1 and 2, find the distance between the points using (a) the Distance Formula and (b) integration.


1. (0, 0), (8, 15) 2. (1, 2), (7, 10)

Finding Arc Length In Exercises 3–16, find the arc length of the graph of the function over the indicated interval.


3. $y = \frac{2}{3}(x^2 + 1)^{3/2}$




4. $y = \frac{x^3}{6} + \frac{1}{2x}$



5. $y = \frac{2}{3}x^{3/2} + 1$



6. $y = 2x^{3/2} + 3$



7. $y = \frac{3}{2}x^{2/3}$, [1, 8]

8. $y = \frac{x^4}{8} + \frac{1}{4x^2}$, [1, 3]

9. $y = \frac{x^5}{10} + \frac{1}{6x^3}$, [2, 5]

10. $y = \frac{3}{2}x^{2/3} + 4$, [1, 27]

11. $y = \ln(\sin x)$, $[\frac{\pi}{4}, \frac{3\pi}{4}]$

12. $y = \ln(\cos x)$, $[0, \frac{\pi}{3}]$

13. $y = \frac{1}{2}(e^x + e^{-x})$, [0, 2]

14. $y = \ln\left(\frac{e^x + 1}{e^x - 1}\right)$, [ln 2, ln 3]

15. $x = \frac{1}{3}(y^2 + 2)^{3/2}$, $0 \leq y \leq 4$

16. $x = \frac{1}{3}\sqrt{y}(y - 3)$, $1 \leq y \leq 4$

Finding Arc Length In Exercises 17–26, (a) sketch the graph of the function, highlighting the part indicated by the given interval, (b) find a definite integral that represents the arc length of the curve over the indicated interval and observe that the integral cannot be evaluated with the techniques studied so far, and (c) use the integration capabilities of a graphing utility to approximate the arc length.

17. $y = 4 - x^2$, $0 \leq x \leq 2$

18. $y = x^2 + x - 2$, $-2 \leq x \leq 1$

19. $y = \frac{1}{x}$, $1 \leq x \leq 3$

20. $y = \frac{1}{x+1}$, $0 \leq x \leq 1$

21. $y = \sin x$, $0 \leq x \leq \pi$

22. $y = \cos x$, $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$

23. $x = e^{-y}$, $0 \leq y \leq 2$

24. $y = \ln x$, $1 \leq x \leq 5$

25. $y = 2 \arctan x$, $0 \leq x \leq 1$

26. $x = \sqrt{36 - y^2}$, $0 \leq y \leq 3$

Approximation In Exercises 27 and 28, determine which value best approximates the length of the arc represented by the integral. (Make your selection on the basis of a sketch of the arc, not by performing any calculations.)

27. $\int_0^2 \sqrt{1 + \left[\frac{d}{dx}\left(\frac{5}{x^2 + 1}\right)\right]^2} dx$

- (a) 25 (b) 5 (c) 2 (d) -4 (e) 3

28. $\int_0^{\pi/4} \sqrt{1 + \left[\frac{d}{dx}(\tan x)\right]^2} dx$

- (a) 3 (b) -2 (c) 4 (d)
- $\frac{4\pi}{3}$
- (e) 1

Approximation In Exercises 29 and 30, approximate the arc length of the graph of the function over the interval [0, 4] in four ways. (a) Use the Distance Formula to find the distance between the endpoints of the arc. (b) Use the Distance Formula to find the lengths of the four line segments connecting the points on the arc when $x = 0$, $x = 1$, $x = 2$, $x = 3$, and $x = 4$. Find the sum of the four lengths. (c) Use Simpson's Rule with $n = 10$ to approximate the integral yielding the indicated arc length. (d) Use the integration capabilities of a graphing utility to approximate the integral yielding the indicated arc length.


29. $f(x) = x^3$

30. $f(x) = (x^2 - 4)^2$


31. Length of a Catenary Electrical wires suspended between two towers form a catenary (see figure) modeled by the equation

$$y = 20 \cosh \frac{x}{20}, \quad -20 \leq x \leq 20$$

where x and y are measured in meters. The towers are 40 meters apart. Find the length of the suspended cable.



32. **Roof Area** A barn is 100 feet long and 40 feet wide (see figure). A cross section of the roof is the inverted catenary $y = 31 - 10(e^{x/20} + e^{-x/20})$. Find the number of square feet of roofing on the barn.



33. **Length of Gateway Arch** The Gateway Arch in St. Louis, Missouri, is modeled by

$$y = 693.8597 - 68.7672 \cosh 0.0100333x, \quad -299.2239 \leq x \leq 299.2239.$$

(See Section 5.8, Section Project: St. Louis Arch.) Use the integration capabilities of a graphing utility to approximate the length of this curve (see figure).




Figure for 33





Figure for 34


34. **Astroid** Find the total length of the graph of the astroid $x^{2/3} + y^{2/3} = 4$.
35. **Arc Length of a Sector of a Circle** Find the arc length from $(0, 3)$ clockwise to $(2, \sqrt{5})$ along the circle $x^2 + y^2 = 9$.
36. **Arc Length of a Sector of a Circle** Find the arc length from $(-3, 4)$ clockwise to $(4, 3)$ along the circle $x^2 + y^2 = 25$. Show that the result is one-fourth the circumference of the circle.

Finding the Area of a Surface of Revolution In Exercises 37–42, set up and evaluate the definite integral for the area of the surface generated by revolving the curve about the x -axis.

37. $y = \frac{1}{3}x^3$



38. $y = 2\sqrt{x}$



39. $y = \frac{x^3}{6} + \frac{1}{2x}, \quad 1 \leq x \leq 2$


40. $y = 3x, \quad 0 \leq x \leq 3$

41. $y = \sqrt{4 - x^2}, \quad -1 \leq x \leq 1$


42. $y = \sqrt{9 - x^2}, \quad -2 \leq x \leq 2$

Finding the Area of a Surface of Revolution In Exercises 43–46, set up and evaluate the definite integral for the area of the surface generated by revolving the curve about the y -axis.

43. $y = \sqrt[3]{x} + 2$



44. $y = 9 - x^2$



45. $y = 1 - \frac{x^2}{4}, \quad 0 \leq x \leq 2$

46. $y = \frac{x}{2} + 3, \quad 1 \leq x \leq 5$

Finding the Area of a Surface of Revolution In Exercises 47 and 48, use the integration capabilities of a graphing utility to approximate the surface area of the solid of revolution.


Function	Interval	Axis of Revolution
47. $y = \sin x$	$[0, \pi]$	x -axis
48. $y = \ln x$	$[1, e]$	y -axis

WRITING ABOUT CONCEPTS


49. **Rectifiable Curve** Define a rectifiable curve.
50. **Precalculus and Calculus** What precalculus formula and representative element are used to develop the integration formula for arc length?
51. **Precalculus and Calculus** What precalculus formula and representative element are used to develop the integration formula for the area of a surface of revolution?



52. HOW DO YOU SEE IT? The graphs of the functions f_1 and f_2 on the interval $[a, b]$ are shown in the figure. The graph of each function is revolved about the x -axis. Which surface of revolution has the greater surface area? Explain.



53. **Think About It** The figure shows the graphs of the functions $y_1 = x$, $y_2 = \frac{1}{2}x^{3/2}$, $y_3 = \frac{1}{4}x^2$, and $y_4 = \frac{1}{8}x^{5/2}$ on the interval $[0, 4]$. To print an enlarged copy of the graph, go to *MathGraphs.com*.



- (a) Label the functions.
- (b) List the functions in order of increasing arc length.
- (c) Verify your answer in part (b) by using the integration capabilities of a graphing utility to approximate each arc length accurate to three decimal places.

54. **Verifying a Formula**

- (a) Given a circular sector with radius L and central angle θ (see figure), show that the area of the sector is given by

$$S = \frac{1}{2} L^2 \theta.$$

- (b) By joining the straight-line edges of the sector in part (a), a right circular cone is formed (see figure) and the lateral surface area of the cone is the same as the area of the sector. Show that the area is $S = \pi rL$, where r is the radius of the base of the cone. (*Hint:* The arc length of the sector equals the circumference of the base of the cone.)




Figure for 54(a)





Figure for 54(b)

- (c) Use the result of part (b) to verify that the formula for the lateral surface area of the frustum of a cone with slant height L and radii r_1 and r_2 (see figure) is $S = \pi(r_1 + r_2)L$. (*Note:* This formula was used to develop the integral for finding the surface area of a surface of revolution.)



55. **Lateral Surface Area of a Cone** A right circular cone is generated by revolving the region bounded by $y = 3x/4$, $y = 3$, and $x = 0$ about the y -axis. Find the lateral surface area of the cone.

56. **Lateral Surface Area of a Cone** A right circular cone is generated by revolving the region bounded by $y = hx/r$, $y = h$, and $x = 0$ about the y -axis. Verify that the lateral surface area of the cone is $S = \pi r \sqrt{r^2 + h^2}$.

57. **Using a Sphere** Find the area of the zone of a sphere formed by revolving the graph of $y = \sqrt{9 - x^2}$, $0 \leq x \leq 2$, about the y -axis.


58. **Using a Sphere** Find the area of the zone of a sphere formed by revolving the graph of $y = \sqrt{r^2 - x^2}$, $0 \leq x \leq a$, about the y -axis. Assume that $a < r$.

59. **Modeling Data** The circumference C (in inches) of a vase is measured at three-inch intervals starting at its base. The measurements are shown in the table, where y is the vertical distance in inches from the base.

y	0	3	6	9	12	15	18
C	50	65.5	70	66	58	51	48

- (a) Use the data to approximate the volume of the vase by summing the volumes of approximating disks.
- (b) Use the data to approximate the outside surface area (excluding the base) of the vase by summing the outside surface areas of approximating frustums of right circular cones.
- (c) Use the regression capabilities of a graphing utility to find a cubic model for the points (y, r) , where $r = C/(2\pi)$. Use the graphing utility to plot the points and graph the model.
- (d) Use the model in part (c) and the integration capabilities of a graphing utility to approximate the volume and outside surface area of the vase. Compare the results with your answers in parts (a) and (b).

60. **Modeling Data** Property bounded by two perpendicular roads and a stream is shown in the figure. All distances are measured in feet.



- (a) Use the regression capabilities of a graphing utility to fit a fourth-degree polynomial to the path of the stream.
- (b) Use the model in part (a) to approximate the area of the property in acres.
- (c) Use the integration capabilities of a graphing utility to find the length of the stream that bounds the property.

61. Volume and Surface Area Let R be the region bounded by $y = 1/x$, the x -axis, $x = 1$, and $x = b$, where $b > 1$. Let D be the solid formed when R is revolved about the x -axis.

- (a) Find the volume V of D .
- (b) Write the surface area S as an integral.
- (c) Show that V approaches a finite limit as $b \rightarrow \infty$.
- (d) Show that $S \rightarrow \infty$ as $b \rightarrow \infty$.

62. Think About It Consider the equation $\frac{x^2}{9} + \frac{y^2}{4} = 1$.

- (a) Use a graphing utility to graph the equation.
- (b) Set up the definite integral for finding the first-quadrant arc length of the graph in part (a).
- (c) Compare the interval of integration in part (b) and the domain of the integrand. Is it possible to evaluate the definite integral? Is it possible to use Simpson's Rule to evaluate the definite integral? Explain. (You will learn how to evaluate this type of integral in Section 8.8.)

Approximating Arc Length or Surface Area In Exercises 63–66, set up the definite integral for finding the indicated arc length or surface area. Then use the integration capabilities of a graphing utility to approximate the arc length or surface area. (You will learn how to evaluate this type of integral in Section 8.8.)

63. Length of Pursuit A fleeing object leaves the origin and moves up the y -axis (see figure). At the same time, a pursuer leaves the point $(1, 0)$ and always moves toward the fleeing object. The pursuer's speed is twice that of the fleeing object. The equation of the path is modeled by

$$y = \frac{1}{3}(x^{3/2} - 3x^{1/2} + 2).$$

How far has the fleeing object traveled when it is caught? Show that the pursuer has traveled twice as far.




Figure for 63




Figure for 64

64. Bulb Design An ornamental light bulb is designed by revolving the graph of

$$y = \frac{1}{3}x^{1/2} - x^{3/2}, \quad 0 \leq x \leq \frac{1}{3},$$

about the x -axis, where x and y are measured in feet (see figure). Find the surface area of the bulb and use the result to approximate the amount of glass needed to make the bulb. (Assume that the glass is 0.015 inch thick.)

65. Astroid Find the area of the surface formed by revolving the portion in the first quadrant of the graph of $x^{2/3} + y^{2/3} = 4$, $0 \leq y \leq 8$, about the y -axis.




Figure for 65




Figure for 66


66. Using a Loop Consider the graph of

$$y^2 = \frac{1}{12}x(4-x)^2$$

shown in the figure. Find the area of the surface formed when the loop of this graph is revolved about the x -axis.

67. Suspension Bridge A cable for a suspension bridge has the shape of a parabola with equation $y = kx^2$. Let h represent the height of the cable from its lowest point to its highest point and let $2w$ represent the total span of the bridge (see figure). Show that the length C of the cable is given by

$$C = 2 \int_0^w \sqrt{1 + (4h^2/w^4)x^2} dx.$$



68. Suspension Bridge The Humber Bridge, located in the United Kingdom and opened in 1981, has a main span of about 1400 meters. Each of its towers has a height of about 155 meters. Use these dimensions, the integral in Exercise 67, and the integration capabilities of a graphing utility to approximate the length of a parabolic cable along the main span.

69. Arc Length and Area Let C be the curve given by $f(x) = \cosh x$ for $0 \leq x \leq t$, where $t > 0$. Show that the arc length of C is equal to the area bounded by C and the x -axis. Identify another curve on the interval $0 \leq x \leq t$ with this property.

PUTNAM EXAM CHALLENGE

70. Find the length of the curve $y^2 = x^3$ from the origin to the point where the tangent makes an angle of 45° with the x -axis.

This problem was composed by the Committee on the Putnam Prize Competition. © The Mathematical Association of America. All rights reserved.