
Range: \_\_\_

9.07b Quiz Review: Centers, Spread, & Normal Distribution

Use the following box and whisker plot to identify the information requested in #1 - 5.



## Notes:

In a normally distributed distribution, the mean is the best measure of central tendency.

With skewed distributions (left or right), however, median can be the better measure of central tendency


The interquartile range (IQR) is the range of values within which reside the middle 50% of the scores.

The first quartile (Q1) indicates that 25% of the scores have a value lower than Q1

The third quartile (Q3) indicates that 75% of the scores have a value lower than Q3

The median is the value that is in the "middle" of the distribution (Q2), with 50% of the scores having a value less than the median

- 6) A machine is used to put bolts into boxes. It does so such that the actual number of bolts in a box is normally distributed with a mean of 106 and a standard deviation of 2.
  - a) Label the x-axis and the Empirical rule probabilities under the Normal distribution graph below.



Use the scenario and graph above to answer the following questions. Show all work and write a probability statement for each. Answers may be given in percent or decimal form.

- a) What percentage of boxes contain no more than 108 bolts?
- b) What percentage of boxes contain at least 104 bolts?
- c) What is the probability of boxes containing between 102 and 112 bolts?\_\_\_\_\_

da en Roy das la carriagnam la Tradición de la como el carrio más la

d) What number represents the 78<sup>h</sup> percentile?

ent jang kasasang sepagai na asah di <mark>Basa neversi</mark>ka tahun 1990 data Kalati

7) A test was given to 120 students, and the scores approximated a normal distribution. If the mean score was 72 with a standard deviation of 7, approximately what percent of the scores were 65 or higher?