

AB PRACTICE TEST II (Larson)
Section I, Part A: Multiple-Choice Questions
 Time: 55 minutes
 Number of Questions: 28


A calculator may not be used on this part of the examination.

1. In what interval is $f(x) = \ln(x^2 - 1)$ decreasing?
 (A) $|x| > 1$
 (B) $|x| \geq 1$
 (C) $x < -1$
 (D) $x > 1$
 (E) $x > 0$

2. Find the limit $\lim_{x \rightarrow 0} \frac{\cos x}{|x|}$.
 (A) 0
 (B) 1
 (C) -1
 (D) π
 (E) The limit does not exist.

3. $\int \left[3^{-x} + \frac{1}{x} \right] dx = ?$
 (A) $3^{-x} + \ln|x| + C$
 (B) $-3^{-x} - \frac{1}{x^2} + C$
 (C) $-3^{-x} \ln 3 + \ln|x| + C$
 (D) $-\frac{3^{-x}}{\ln 3} + \ln|x| + C$
 (E) $-\frac{3^{-x}}{\ln 3} - \frac{1}{x^2} + C$

4. Find $\frac{dy}{dx}$ for $\cos(x+y) = x$.
 (A) $-\csc(x+y) - 1$
 (B) $\frac{\cos(x+y)}{\sin^2(x+y)}$
 (C) $\frac{x}{1-x^2}$
 (D) $-\sin(x+y) \cdot \cot(x+y)$
 (E) $-\sin(x+y) - 1$



The graph of $f(x)$ consists of four line segments and a semicircle as shown above in the closed interval $-3 \leq x \leq 5$. Let g be the function given by $g(x) = \int_0^x f(t) dt$. Use this information for problems 5-7.

5. What is $g(-1) + g'(-1) + g''(-1)$?
 (A) -1
 (B) 0
 (C) 1
 (D) 2
 (E) 3
6. What is $\int_{-3}^5 f(t) dt$?
 (A) $7 - \pi$
 (B) $7 - \frac{\pi}{2}$
 (C) $7 - \frac{\pi}{4}$
 (D) $12 - \frac{\pi}{2}$
 (E) $12 - \frac{\pi}{4}$

7. Which of the following statements is false for $g(x)$?
- (A) The absolute maximum for $g(x)$ occurs at $x = 5$.
 - (B) A relative minimum for $g(x)$ occurs at $x = -1$.
 - (C) A point of inflection for $g(x)$ occurs at $x = 3$.
 - (D) $g(x)$ has roots at $x = 0$ and $x = -2$.
 - (E) $g(x)$ is concave down in the open interval $-2 < x < -1$.
8. The position of a particle moving along the x -axis is given by $x(t) = 2 + 3t - t^3$. What is the speed of the particle at $t = 4$?
- (A) -50
 - (B) -45
 - (C) 32
 - (D) 45
 - (E) 50
9. A region R is bounded by the curve $x = y^2 - 1$ and the y -axis. What is the volume generated when region R is rotated about the y -axis?
- (A) $\frac{\pi}{2}$
 - (B) $\frac{8\pi}{15}$
 - (C) π
 - (D) $\frac{16\pi}{15}$
 - (E) $\frac{4\pi}{3}$
10. Which of the following statements is true for $f(x) = \sqrt[3]{x} + 1$?
- I. $f(x)$ is always increasing, $x \neq 0$.
 - II. The tangent to the curve at $x = 0$ is horizontal.
 - III. The Mean Value Theorem can be applied to $f(x)$ in the closed interval $-1 \leq x \leq 1$.
- (A) I only
 - (B) II only
 - (C) III only
 - (D) II and III only
 - (E) I, II, and III

11. The acceleration of a model car along an incline is given by

$$a(t) = \frac{t^2 + t}{t^2 + 1} \text{ cm/sec}^2, \text{ for } 0 \leq t < 1. \text{ If}$$


$$v(0) = 1 \text{ cm/sec, what is } v(t)?$$

- (A) $\tan^{-1} t + \frac{1}{2} \ln(t^2 + 1) + 1 \text{ cm/sec}$
- (B) $\tan^{-1} t - \frac{1}{2} \ln(t^2 + 1) + 1 \text{ cm/sec}$
- (C) $t - \frac{1}{2} \ln(t^2 + 1) - \tan^{-1} t + 1 \text{ cm/sec}$
- (D) $t + \frac{1}{2} \ln(t^2 + 1) + \tan^{-1} t + 1 \text{ cm/sec}$
- (E) $t - \frac{1}{2} \ln(t^2 + 1) + \tan^{-1} t + 1 \text{ cm/sec}$

t	$R(t)$
0	12
2	18
4	10
6	15
8	12
10	16
12	8

12. Water is dripping into a vase at a variable rate. The rate, $R(t)$ in cm^3/min , is recorded every 2 mins for 12 mins, as listed in the chart above. Using a right Riemann sum with 3 equal intervals, find the approximate average rate at which the water drips into the vase over the 12 mins.
- (A) $10 \text{ cm}^3/\text{min}$
 - (B) $10\frac{1}{3} \text{ cm}^3/\text{min}$
 - (C) $16\frac{1}{3} \text{ cm}^3/\text{min}$
 - (D) $40 \text{ cm}^3/\text{min}$
 - (E) $41\frac{1}{3} \text{ cm}^3/\text{min}$

13. If $h''(x) = e^{x-1}(2x-1)^2(x-3)^3(4x+5)$, then $h(x)$ has how many points of inflection?
- (A) 4
(B) 3
(C) 2
(D) 1
(E) 0




14. The graph of $f'(x)$ is shown in the figure above. If $\int_1^2 f'(x) dx = k$, what is $f(2) - f(-1)$?
- (A) $k + \frac{11}{6}$
(B) $k + \frac{3}{2}$
(C) $k - \frac{1}{2}$
(D) $k - 1$
(E) $k - \frac{4}{3}$

15. $\int \frac{1}{16+x^2} dx =$
- (A) $4 \tan^{-1} x + C$
(B) $\frac{1}{4} \tan^{-1} x + C$
(C) $\tan^{-1} \frac{x}{4} + C$
(D) $\frac{1}{4} \tan^{-1} 4x + C$
(E) $\frac{1}{4} \tan^{-1} \frac{x}{4} + C$

16. A particle moves along the x -axis so that its velocity for $t \geq 0$ is given by $v(t) = -t^3 + \frac{9}{4}t^4$. At what time is the acceleration of the particle a minimum?

- (A) $\frac{2}{9}$
(B) $\frac{1}{3}$
(C) $\frac{4}{9}$
(D) 0
(E) none of these
17. What is the value of $\lim_{h \rightarrow 0} \frac{\sin^{-1}(1+h) - \frac{\pi}{2}}{h}$?
- (A) 1
(B) 0
(C) -1
(D) $\frac{\pi}{2}$
(E) The limit does not exist.

18. Which of the following is the equation of the tangent to the curve of $f(x) = e^{\sin x} + x$ at $x = 0$?
- (A) $y = 2x - 1$
(B) $y = 2x + 1$
(C) $y = 2x$
(D) $y = 1$
(E) $y = 0$



19. Which of the following statements matches the slope field shown above?

- (A) $\frac{dy}{dx} = \frac{x-y}{2y}$
 (B) $\frac{dy}{dx} = \frac{x+y}{2y}$
 (C) $\frac{dy}{dx} = \frac{x-y}{2x}$
 (D) $\frac{dy}{dx} = \frac{x+y}{2x}$
 (E) $\frac{dy}{dx} = \frac{2y}{x-y}$

20. If $F(x) = \int_{3x-2}^2 f(2t) dt$, what is $F'(x)$

- in terms of x ?
 (A) $3f(6x-4)$
 (B) $f(3x-2)$
 (C) $\frac{1}{2}f(3x-2)$
 (D) $-f(3x-2)$
 (E) $-3f(6x-4)$

21. What is the slope of the line normal to the curve $h(x) = \sqrt{5x^3 - 2x + 1}$ at the point where $x = 1$?

- (A) $-\frac{13}{4}$
 (B) $-\frac{4}{13}$
 (C) $\frac{4}{13}$
 (D) $\frac{13}{4}$
 (E) none of these

22. Let $y = 2x(\sin 2x + x \cos 2x)$ in the

interval $0 \leq x \leq \frac{\pi}{2}$. What is the average rate of change of y with respect to x in this interval?

- (A) $-\pi$
 (B) $-\frac{\pi}{2}$
 (C) 0
 (D) $\frac{\pi}{2}$
 (E) π

23. Find k so that the relative minimum of $f(x) = x^3 - \ln(k+x)$ occurs at $x = 1$.

- (A) 2
 (B) $\sqrt{3}$
 (C) $\frac{1}{3}$
 (D) 0
 (E) $-\frac{2}{3}$

24. The concentration of an anti-inflammatory drug in the bloodstream t mins after taking a

single dose is $C(t) = \frac{2t}{8100 + t^2}$, $t \geq 0$.

At what time is the concentration the greatest?

- (A) 90 minutes
 (B) $30\sqrt{6}$ minutes
 (C) $30\sqrt{3}$ minutes
 (D) $15\sqrt{6}$ minutes
 (E) none of these

25. If $\lim_{x \rightarrow \infty} \frac{ae^x}{x + e^x} = 3$, what is a ?

- (A) 9
(B) 6
(C) 3
(D) none of these
(E) No such value of a exists.

x	$f(x)$	$g(x)$	$f'(x)$	$g'(x)$
1	3	4	$\frac{2}{3}$	$-\frac{5}{2}$
2	4	2	$\frac{4}{3}$	$-\frac{3}{2}$
4	8	1	$\frac{8}{3}$	$-\frac{1}{2}$

26. Use the values listed in the chart above to find the value of

$$\frac{d}{dx} [f(g(x^2))] \text{ when } x = 2.$$

- (A) -8
(B) -4
(C) $-\frac{4}{3}$
(D) $\frac{2}{3}$
(E) $\frac{8}{3}$

27. The volume of an open rectangular box is 8 cm^3 , and the length of the rectangular base is twice as long as its width. What is the width of the base so that the surface area of the open box is minimized?

- (A) $\sqrt[3]{3}$
(B) $\sqrt[3]{6}$
(C) $\sqrt{3}$
(D) 2
(E) $\sqrt{6}$

28. The base of a solid is bounded by the curve $y = \sqrt{x+1}$, the x -axis, and the line $x = 1$. The cross sections, taken perpendicular to the x -axis, are squares. Find the volume of the solid.

- (A) $\frac{1}{2}$
(B) 1
(C) $\frac{4\sqrt{2}}{3}$
(D) 2
(E) $\frac{8}{3}$

Section I, Part B: Multiple-Choice Questions

Time: 50 minutes

Number of Questions: 17

A calculator may be used on this part of the examination.

29. Suppose $g(0) = 4$, $g'(0) = 8$, and $g''(0) = -12$. If $h(x) = \sqrt{g(x)}$, what is

- $h''(0)$?
- (A) -5
(B) $-\frac{13}{4}$
(C) $-\frac{1}{32}$
(D) $\frac{3}{8}$
(E) 1

30. If $g''(x) = \frac{x}{2+e^x}$ and $g'(0) = -1$, what is $g'(3)$?

- (A) -0.864
(B) -0.473
(C) 0.136
(D) 0.527
(E) 1.527

31. Two cars are converging on a point P as they drive at right angles to each other. Car A is traveling at 60 miles per hour and car B is traveling at 50 miles per hour. At the instant when car A is 12 miles from the point P and car B is 10 miles from the point P , at what rate is the distance between the cars decreasing?

(A) 55 miles per hour
(B) 55.455 miles per hour
(C) 76.882 miles per hour
(D) 78.102 miles per hour
(E) none of these

32. A line tangent to the curve

$f(x) = \frac{1}{2^{2x}}$ at the point $(a, f(a))$ has a slope of -1 . What is the x -intercept of this tangent?

(A) 0.236
(B) 0.500
(C) 0.721
(D) 0.957
(E) 1.000


33. Consider the function $g(x) = \tan(x+2)$ in the open interval $-4 < x < 5$. How many times are the tangents to $g(x)$ parallel to the line $y = 2x - 1$?

(A) never
(B) 2
(C) 4
(D) 5
(E) an infinite number of times





34. What is the sum of all k values that

satisfy $\int_1^{2k} x - \frac{k}{x^2} dx = 15$?

(A) 3
(B) $\frac{1}{2}$
(C) 1
(D) 0
(E) $-\frac{1}{2}$



35. The graph of $f'(x)$ is shown above. Which of the following could be a graph of $f(x)$?



(E) none of these

t	0	1	2	3	4
$H(t)$	0	1.3	1.5	2.1	2.6

36. A small plant is purchased from a nursery and the change in the height of the plant is measured at the end of every day for four days. The change in the height of the plant is listed in the chart above where $H(t)$ is in inches per day and t is in days. Using the Trapezoidal Rule, which of the following represents an estimate of the average rate of growth of the plant during the 4-day period?

- (A) $\frac{1}{4}(0+1.3+1.5+2.1+2.6)$
 (B) $\frac{1}{4}\left[\frac{1}{2}(0+1.3+1.5+2.1+2.6)\right]$
 (C) $\frac{1}{4}\left\{\frac{1}{2}[0+2(1.3)+2(1.5)+2(2.1)+2.6]\right\}$
 (D) $\frac{1}{4}\left\{\frac{1}{2}[0+2(1.3)+2(1.5)+2(2.1)+2(2.6)]\right\}$
 (E) $\frac{1}{4}\left\{\frac{1}{4}[0+2(1.3)+2(1.5)+2(2.1)+2.6]\right\}$


37. A bug travels along the y -axis such that its velocity, in centimeters per second, is given by $v(t) = (t^2 - 1)e^{t+2t}$, and t is in seconds. How far does the bug travel in the first 1.5 sec?

- (A) 2.257 cm
 (B) 3.386 cm
 (C) 27.155 cm
 (D) 29.412 cm
 (E) 31.669 cm

38. Consider a differentiable function $f(x)$ that has an x -intercept of 2 and a y -intercept of 4. In the interval $0 < x < 2$, $f(x)$ is decreasing and concave down. Which of the following must be true?

- I. The Mean Value Theorem is satisfied when $c = 1$.
 II. $\int_0^1 f(x) dx > \int_1^2 f(x) dx$
 III. A tangent approximation of the function for any value in the interval $0 < x < 2$ will underestimate the function value.

- (A) I only
 (B) II only
 (C) III only
 (D) I and II only
 (E) II and III only



Let $f(x) = x^2 + \int_{-2}^x g(t) dt$, where $g(x)$ is shown in the graph above. Use this graph to answer problems 39–41.

39. What is $f(-2)$?

- (A) -6
 (B) -4
 (C) 0
 (D) 2
 (E) 4

40. What is $f'(-2)$?

- (A) -6
 (B) -4
 (C) 0
 (D) 2
 (E) 4

41. What is $f''(2)$?
- (A) -6
(B) -4
(C) 0
(D) 2
(E) 4
42. What is the area of the regions between $f(x) = 2x \sin(2x)$ and $g(x) = -2x \cos 2x$ on the interval $0 \leq x \leq \frac{\pi}{2}$?
- (A) 0.571
(B) 0.595
(C) 1.166
(D) 1.178
(E) 1.761
43. The graph of $f'(x)$ is given above, in the interval $-8 \leq x \leq 5$. In which interval(s) is $f(x)$ decreasing and concave down?
- (A) $-2 < x < 0$
(B) $-2 < x < 3$ and $3 < x < 5$
(C) $-2 < x < 0$ and $3 < x < 5$
(D) $-6 < x < 0$ and $3 < x < 5$
(E) $-8 < x < -3$ and $2 < x < 5$
44. The rate of decay of a radioactive isotope is directly proportional to the amount remaining. If the half-life of the radioactive isotope, Einsteinium, is 276 days and a sample initially weighs 25 grams, what is its rate of decay on the 120th day?
- (A) -0.046 grams per day
(B) -0.031 grams per day
(C) -0.003 grams per day
(D) -0.002 grams per day
(E) -0.001 grams per day
45. A region is bounded by the function $y = \sin^{-1}(x-1) + \frac{\pi}{2}$, the line $x = 2$, and the x -axis. A solid is formed when the region is rotated about the line $x = 2$. What is the radius of the volume of rotation?
- (A) $r = \sin^{-1}(x-1) + \frac{\pi}{2}$
(B) $r = 2 - \left(\sin^{-1}(x-1) + \frac{\pi}{2} \right)$
(C) $r = 2 - \sin\left(y - \frac{\pi}{2}\right)$
(D) $r = 1 - \sin\left(y - \frac{\pi}{2}\right)$
(E) $r = 1 + \sin\left(y - \frac{\pi}{2}\right)$

Answers and Answer Explanations

Using the table below, score your test. Determine how many questions you answered correctly and how many you answered incorrectly. Additional information about scoring is at the end of the Practice Test.

1. C	2. E	3. D	4. A	5. C
6. B	7. E	8. D	9. D	10. A
11. E	12. A	13. C	14. C	15. E
16. A	17. E	18. B	19. A	20. E
21. B	22. A	23. E	24. A	25. C
26. C	27. B	28. D	29. A	30. B
31. D	32. D	33. D	34. B	35. D
36. C	37. E	38. B	39. E	40. A
41. D	42. E	43. C	44. A	45. D

MULTIPLE-CHOICE QUESTIONS

1. ANSWER: (C) The domain for $f(x) = \ln(x^2 - 1)$ is $|x| > 1$. With $f'(x) = \frac{2x}{x^2 - 1} = 0 \Rightarrow x = 0$, but $x = 0$ is not in the domain of $f(x)$. The behavior of the curve is analyzed using the sign chart below.

x	$-\infty < x < -1$	$1 < x < \infty$
$f'(x)$	negative	positive
$f(x)$	decreasing	increasing

Therefore the curve is decreasing when $x < -1$.
(Calculus 7th ed. pages 174–180 / 8th ed. pages 179–185)

2. ANSWER: (E) $\lim_{x \rightarrow 0} \frac{\cos x}{|x|} = \frac{1}{0}$, which does not exist. (The value of the limit from both the left and right side of 0 is ∞ .)
(Calculus 7th ed. pages 80–84 / 8th ed. pages 83–87)

3. ANSWER: (D) By substituting $u = -x$ and $du = -dx$,
$$\int 3^{-x} + \frac{1}{x} dx = -\int 3^u du + \int \frac{1}{x} dx = -\frac{3^u}{\ln 3} + \ln |x| + C$$

(Calculus 7th ed. pages 351–356, 324–330 / 8th ed. pages 360–365, 332–337)