

## BC Calculus – 9.1 Notes – Defining and Differentiating Parametric Equations

We have been looking at graphs of one equation with two variables, typically  $x$  and  $y$ . Now we are looking at three variables that will represent a curve in the plane.


In the rectangular equation we are able to determine where the object is located at a point  $(x, y)$ , but with the addition of the third variable (often  $t$ ), we are able to determine when the object was at a point  $(x, y)$ . NOTE: the third variable  $t$  is often time, but not always.

### Parametric Equations

If  $f$  and  $g$  are continuous functions of  $t$  on an interval  $I$ , then the equations  $x = f(t)$  and  $y = g(t)$  are parametric equations and  $t$  is the parameter. You can sketch the curve of a parametric by substituting in values for  $t$ .

- Sketch the curve with the following parametrization:  $x(t) = 2t$  and  $y(t) = t^2 - 1$ , with  $-1 \leq t \leq 2$ .

$t$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$	2
$x$							
$y$							



```

WINDOW
Tmin=-1
Tmax=2
Tstep=0.2
Xmin=-3.5
Xmax=5
Xscl=1
Ymin=-2
Ymax=4.5
Yscl=1
    
```

To find the rectangular equation when you are given the parametric equations, eliminate the parameter  $t$  through substitution.

- Given  $x(t) = 2t, y(t) = t^2 - 1$ . Find the rectangular equation by eliminating the parameter.

- Given the parametric equations  $x(t) = 2 \cos t$  and  $y(t) = 2 \sin t$ . Eliminate the parameter.

2

### Derivative of a Parametric Equation

The derivative of a parametric given by  $x = f(t)$  and  $y = g(t)$  is found by the following:

4. Given  $x(t) = t^{\frac{1}{2}}$  and  $y(t) = \frac{1}{4}(t^2 - 4)$  for  $t \geq 0$ . Find  $\frac{dy}{dx}$

5. Given  $x(t) = e^{2t}$  and  $y(t) = \cos t$  for  $t \geq -1$ . Find the equation of a tangent line when  $t = \frac{\pi}{2}$ .

#### 9.1 Practice Problems:

1. For the given parametric equations, eliminate the parameter and write the corresponding rectangular equation.  $x = e^{-t}$  and  $y = e^{2t} - 1$ .

2. Let  $C$  be a curve described by the parametrization  $x = 5t$  and  $y = t^4 + 3$ . Find an expression for the slope of the line tangent to  $C$  at any point  $(x, y)$ .

3. The position of a particle at any time  $t \geq 0$  is given by  $x(t) = 3t^2 + 1$  and  $y(t) = \frac{2}{3}t^3$ . Find  $\frac{dy}{dx}$  as a function of  $x$ .

4. A particle moves along the curve  $xy + y = 9$ . If  $x = 2$  and  $\frac{dy}{dt} = 3$ , what is the value of  $\frac{dx}{dt}$ ?

5. A curve is described by the parametric equations  $x = t \cos t$  and  $y = t \sin t$ . Find the equation of the line tangent to the curve at the point determined by  $t = \pi$ .
6. **Calculator active.** The coordinates  $(x(t), y(t))$  of the position of a drone change at rates given by  $x'(t) = 2t^3$  and  $y'(t) = t^{\frac{1}{2}}$ , where  $x(t)$  and  $y(t)$  are measured in meters and  $t$  is measured in seconds. At what time  $t$ , for  $0 \leq t \leq 2$ , does the slope of the line tangent to its path have a slope of 1.5?
- 
7. A curve in the  $xy$ -plane is defined by the parametric equations  $x(t) = \cos(3t)$  and  $y(t) = \sin(3t)$  for  $t \geq 0$ . What is the value of  $\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}$ ?
8. A curve is defined by the parametric equations  $x(t) = at^2 + b$  and  $y(t) = ct - b$ , where  $a$ ,  $b$ , and  $c$  are nonzero constants. What is the slope of the line tangent to the curve at the point  $(x(t), y(t))$  when  $t = 2$ ?
- 
9. **No Calculator.** For  $0 \leq t \leq 11$  the parametric equations  $x = 3 \sin t$  and  $y = 2 \cos t$  describe the elliptical path of an object. At the point where  $t = 11$ , the object travels along a line tangent to the path at that point. What is the slope of that line?
10. A particle moves in the  $xy$ -plane so that its position for  $t \geq 0$  is given by the parametric equations  $x(t) = 2kt^2$  and  $y(t) = 3t$ , where  $k$  is a positive constant. When  $t = 2$  the line tangent to the particle's path has a slope of 4. What is the value of  $k$ ?

4

11. Find the equation of the line tangent to the curve defined parametrically by the equations  $x(t) = t^3 + 2t$  and  $y(t) = 2t^4 + 2t^2$  when  $t = 1$ .
12. For what values of  $t$  does the curve given by the parametric equations  $x(t) = \frac{1}{4}t^4 - \frac{9}{2}t^2$  and  $y(t) = 3t^3 + 2t$  have a vertical tangent?

- 
13. Suppose a curve is given by the parametric equations  $x = f(t)$  and  $y = g(t)$ , for all  $t > 1$  and  $\frac{dy}{dt} = \frac{t^2+2}{t-1} * \frac{dx}{dt}$ . What is the value of  $\frac{dy}{dx}$  when  $t = 2$ ?

## 9.1 Parametric Equations

## Test Prep

14. A curve is defined parametrically by  $x(t) = t^2$  and  $y(t) = t^3 - 3t$ . Find the points on the graph where the tangent line is horizontal or vertical.

15. **Free Response.** Consider the curve given by the parametric equations  $y = t^3 - 12t$  and  $x = \frac{1}{2}t^2 - t$ .

a. Find  $\frac{dy}{dx}$  in terms of  $t$ .

b. Write an equation for the line tangent to the curve at the point where  $t = -1$ .

c. Find the  $x$  and  $y$  coordinates for each critical point on the curve and identify each point as having a vertical or horizontal tangent.

---

16. A curve is given by the parametric equations  $x(t) = 5t^3 - 5$  and  $y(t) = t^2 + 7$ . What is the equation of the tangent line to the curve when  $t = 1$ ?

A.  $x = 0$

B.  $y = \frac{2}{15}x + 8$

C.  $y = \frac{2}{15}x + 1$

D.  $y = 8$

E.  $y = \frac{15}{2}x + 7$

6

9.1 AP Practice Problems (p.651) – Parametric Equations

1. Which pair of parametric equations represents a plane curve that is a circle with a radius of 3 and a center at  $(0, 0)$ ?

(A)  $x(t) = 9 \cos t, y(t) = 9 \sin t, 0 \leq t \leq 2\pi$

(B)  $x(t) = 3 \sin t, y(t) = 3 \cos t, 0 \leq t \leq \pi$

(C)  $x(t) = 3 \sin(2t), y(t) = 3 \cos(2t), 0 \leq t \leq \pi$

(D)  $x(t) = \cos(3t), y(t) = \sin(3t), 0 \leq t \leq 2\pi$

2. A rectangular equation of the curve whose parametric equations are  $x(t) = t + 1, y(t) = t^2 + 3t$  is

(A)  $y = x^2 + x - 2$       (B)  $y = x^2 + 3x$

(C)  $y = x^2 + 5x + 4$       (D)  $y = x^2 + 3x - 2$

3. A rectangular equation of the parametric equations  $x(t) = 4 \cos t$ ,  $y(t) = \frac{1}{2} \sin t$  is

(A)  $\frac{x^2}{16} + \frac{y^2}{4} = 1$       (B)  $\frac{x^2}{16} + 2y^2 = 1$

(C)  $16x^2 + 4y^2 = 1$       (D)  $\frac{x^2}{16} + 4y^2 = 1$

4. An object is moving along a plane curve according to the parametric equations

$$x(t) = -5 \cos\left(\frac{\pi}{2}t\right), y(t) = 2 \sin\left(\frac{\pi}{2}t\right), 0 \leq t \leq 8$$

- (a) Find a rectangular equation for the parametric equations.
- (b) Describe the motion of the object from  $t = 0$  to  $t = 8$ . Be sure to include where the object begins, where it ends, its path, and its orientation.