


BC Calculus – 9.4a Notes – Defining and Differentiating in Polar Form

(x, y) is for a rectangular coordinate system.

(r, θ) is for a polar coordinate system.

r is a directed distance from the origin to a point P.

θ is the directed angle



| Polar \iff Rectangular | Rectangular \iff Polar |
|--------------------------|-----------------------------|
| $x = r \cos \theta$ | $\tan \theta = \frac{y}{x}$ |
| $y = r \sin \theta$ | $r^2 = x^2 + y^2$ |

Convert the following from polar form to rectangular form.

1. $r \cos \theta = -4$

2. $4r \cos \theta = r^2$

3. $\frac{4}{2 \cos \theta - \sin \theta} = r$

Slope of a Curve in Polar Form


A curve in polar form is given by $r = f(\theta)$, then its rectangular coordinates are given by $\begin{cases} x = f(\theta) \cos \theta \\ y = f(\theta) \sin \theta \end{cases}$. The derivative $\frac{dy}{dx}$ is defined the same way as the derivative of a parametric equation.

$$\frac{dy}{dx} =$$

The following is an example of a common problem found on the AP Exam!

4. What is the slope of the line tangent to the polar curve $r = 1 + 2 \sin \theta$ at $\theta = 0$?

5. Find the value(s) of θ where the polar graph $r = 1 - \sin \theta$ on the interval $0 \leq \theta \leq 2\pi$ has horizontal and vertical tangent lines.




Practice Problems

Problems 1-5 are pre-calculus review on polar form.

- | | |
|--|--|
| <p>1. Find the corresponding rectangular coordinates for the polar coordinates $(7, \frac{5\pi}{4})$.</p> | <p>2. Calculator active. Find two sets of polar coordinates for the rectangular coordinate $(4, -2)$. Limit your answers on the interval $0 \leq \theta \leq 2\pi$.</p> |
| <p>3. Convert the rectangular equation $x^2 + y^2 = 16$ to a polar equation.</p> | <p>4. Convert the polar equation $r = 3 \sec \theta$ to a rectangular equation.</p> |

5. Sketch the polar curve $r = 2 \cos 3\theta$ for $0 \leq \theta \leq \pi$ **without** a calculator, then check your answer.



Find the slope of the line tangent to the polar curve at the given value of θ .

6. $r = 3\theta$ at $\theta = \frac{\pi}{2}$.

7. $r = \frac{5}{3 - \cos \theta}$ at $\theta = \frac{3\pi}{2}$.

8. $r = \cos \theta$ at $\theta = \frac{\pi}{3}$.

9. $r = 2(1 - \sin \theta)$ at $\theta = 0$.

10. A particle moves along the polar curve $r = 3 \cos \theta$ so that $\frac{d\theta}{dt} = 2$. Find the value of $\frac{dr}{dt}$ at $\theta = \frac{\pi}{3}$. *Hint: remember implicit differentiation?*
11. A polar curve is given by the equation $r = \frac{15\theta}{\theta^2+1}$ for $\theta \geq 0$. What is the instantaneous rate of change of r with respect to θ when $\theta = 1$?
-
12. Find the value(s) of θ where the polar graph $r = 2 - 2 \cos \theta$ has a horizontal tangent line on the interval $0 \leq \theta \leq 2\pi$. Use a graphing calculator to verify your answers.
13. Find the value(s) of θ where the polar graph $r = 3 - 3 \sin \theta$ has a vertical tangent line on the interval $0 \leq \theta \leq 2\pi$. Use a graphing calculator to verify your answers.
-
14. **Calculator active.** For a certain polar curve $r = f(\theta)$, it is known that $\frac{dx}{d\theta} = \cos \theta - \theta \sin \theta$ and $\frac{dy}{d\theta} = \sin \theta + \theta \cos \theta$. What is the value of $\frac{d^2y}{dx^2}$ at $\theta = 3$?

9.4a Differentiating in Polar Form

15. A polar curve is given by the differentiable function $r = f(\theta)$ for $0 \leq \theta \leq 2\pi$. If the line tangent to the polar curve at $\theta = \frac{\pi}{6}$ is vertical, which of the following must be true?

- A. $f\left(\frac{\pi}{6}\right) = 0$ B. $f'\left(\frac{\pi}{6}\right) = 0$ C. $\frac{1}{2}f\left(\frac{\pi}{6}\right) - \frac{\sqrt{3}}{2}f'\left(\frac{\pi}{6}\right) = 0$ D. $\frac{\sqrt{3}}{2}f'\left(\frac{\pi}{6}\right) - \frac{1}{2}f\left(\frac{\pi}{6}\right) = 0$

16. **Calculator active.** For $0 \leq t \leq 8$, a particle moving in the xy -plane has position vector $\langle x(t), y(t) \rangle = \langle \sin(2t), t^2 - t \rangle$, where $x(t)$ and $y(t)$ are measured in meters and t is measured in seconds.

- a. Find the speed of the particle at time $t = 3$ seconds. Indicate units of measure.
- b. At time $t = 5$ seconds, is the speed of the particle increasing or decreasing? Explain your answer.
- c. Find the total distance the particle travels over the time interval $0 \leq t \leq 6$ seconds.
- d. At time $t = 8$ seconds, the particle begins moving in a straight line. For $t \geq 8$, the particle travels with the same velocity vector that it had at time $t = 8$ seconds. Find the position of the particle at time $t = 11$ seconds.