


1)

The figure above shows the graph of f' , the derivative of a twice-differentiable function f , on the interval $[-3, 4]$. The graph of f' has horizontal tangents at $x = -1$, $x = 1$, and $x = 3$. The areas of the regions bounded by the x -axis and the graph of f' on the intervals $[-2, 1]$ and $[1, 4]$ are 9 and 12, respectively.



- (a) Find all x -coordinates at which f has a relative maximum. Give a reason for your answer.
- (b) On what open intervals contained in $-3 < x < 4$ is the graph of f both concave down and decreasing? Give a reason for your answer.
- (c) Find the x -coordinates of all points of inflection for the graph of f . Give a reason for your answer.
- (d) Given that $f(1) = 3$, write an expression for $f(x)$ that involves an integral. Find $f(4)$ and $f(-2)$.

(a) $f'(x) = 0$ at $x = -2$, $x = 1$, and $x = 4$.
 $f'(x)$ changes from positive to negative at $x = -2$.
 Therefore, f has a relative maximum at $x = -2$.

2 : { 1 : identifies $x = -2$
 1 : answer with reason

(b) The graph of f is concave down and decreasing on the intervals $-2 < x < -1$ and $1 < x < 3$ because f' is decreasing and negative on these intervals.

2 : { 1 : intervals
 1 : reason

(c) The graph of f has a point of inflection at $x = -1$ and $x = 3$ because f' changes from decreasing to increasing at these points.

2 : { 1 : identifies $x = -1, 1, \text{ and } 3$
 1 : reason

The graph of f has a point of inflection at $x = 1$ because f' changes from increasing to decreasing at this point.

(d) $f(x) = 3 + \int_1^x f'(t) dt$

$$f(4) = 3 + \int_1^4 f'(t) dt = 3 + (-12) = -9$$

$$f(-2) = 3 + \int_1^{-2} f'(t) dt = 3 - \int_{-2}^1 f'(t) dt = 3 - (-9) = 12$$


3 : { 1 : integrand
 1 : expression for $f(x)$
 1 : $f(4)$ and $f(-2)$

2)

The figure above shows the graph of the piecewise-linear function f . For $-4 \leq x \leq 12$, the function g is defined by

$$g(x) = \int_2^x f(t) dt.$$

- (a) Does g have a relative minimum, a relative maximum, or neither at $x = 10$? Justify your answer.
- (b) Does the graph of g have a point of inflection at $x = 4$? Justify your answer.
- (c) Find the absolute minimum value and the absolute maximum value of g on the interval $-4 \leq x \leq 12$. Justify your answers.
- (d) For $-4 \leq x \leq 12$, find all intervals for which $g(x) \leq 0$.



Graph of f

- (a) The function g has neither a relative minimum nor a relative maximum at $x = 10$ since $g'(x) = f(x)$ and $f(x) \leq 0$ for $8 \leq x \leq 12$.
- (b) The graph of g has a point of inflection at $x = 4$ since $g'(x) = f(x)$ is increasing for $2 \leq x \leq 4$ and decreasing for $4 \leq x \leq 8$.
- (c) $g'(x) = f(x)$ changes sign only at $x = -2$ and $x = 6$.

x	$g(x)$
-4	-4
-2	-8
6	8
12	-4

On the interval $-4 \leq x \leq 12$, the absolute minimum value is $g(-2) = -8$ and the absolute maximum value is $g(6) = 8$.

- (d) $g(x) \leq 0$ for $-4 \leq x \leq 2$ and $10 \leq x \leq 12$.

1 : $g'(x) = f(x)$ in (a), (b), (c), or (d)

1 : answer with justification

1 : answer with justification

4 : { 1 : considers $x = -2$ and $x = 6$
as candidates
1 : considers $x = -4$ and $x = 12$
2 : answers with justification

2 : intervals

3)

$$\begin{aligned}
 \text{(a)} \quad f(-5) &= f(1) + \int_1^{-5} g(x) \, dx = f(1) - \int_{-5}^1 g(x) \, dx \\
 &= 3 - \left(-9 - \frac{3}{2} + 1\right) = 3 - \left(-\frac{19}{2}\right) = \frac{25}{2}
 \end{aligned}$$

$$2: \begin{cases} 1: \text{integral} \\ 1: \text{answer} \end{cases}$$

$$\begin{aligned}
 \text{(b)} \quad \int_1^6 g(x) \, dx &= \int_1^3 g(x) \, dx + \int_3^6 g(x) \, dx \\
 &= \int_1^3 2 \, dx + \int_3^6 2(x-4)^2 \, dx \\
 &= 4 + \left[\frac{2}{3}(x-4)^3\right]_{x=3}^{x=6} = 4 + \frac{16}{3} - \left(-\frac{2}{3}\right) = 10
 \end{aligned}$$

$$3: \begin{cases} 1: \text{split at } x = 3 \\ 1: \text{antiderivative of } 2(x-4)^2 \\ 1: \text{answer} \end{cases}$$

(c) The graph of f is increasing and concave up on $0 < x < 1$ and $4 < x < 6$ because $f'(x) = g(x) > 0$ and $f''(x) = g'(x)$ is increasing on those intervals.

$$2: \begin{cases} 1: \text{intervals} \\ 1: \text{reason} \end{cases}$$

(d) The graph of f has a point of inflection at $x = 4$ because $f'(x) = g(x)$ changes from decreasing to increasing at $x = 4$.

$$2: \begin{cases} 1: \text{answer} \\ 1: \text{reason} \end{cases}$$

4)

(a) $f(-6) = f(-2) + \int_{-2}^{-6} f'(x) dx = 7 - \int_{-6}^{-2} f'(x) dx = 7 - 4 = 3$
 $f(5) = f(-2) + \int_{-2}^5 f'(x) dx = 7 - 2\pi + 3 = 10 - 2\pi$

3 : $\begin{cases} 1 : \text{uses initial condition} \\ 1 : f(-6) \\ 1 : f(5) \end{cases}$

(b) $f'(x) > 0$ on the intervals $[-6, -2]$ and $(2, 5)$.
 Therefore, f is increasing on the intervals $[-6, -2]$ and $[2, 5]$.

2 : answer with justification

(c) The absolute minimum will occur at a critical point where $f'(x) = 0$ or at an endpoint.

2 : $\begin{cases} 1 : \text{considers } x = 2 \\ 1 : \text{answer with justification} \end{cases}$

$f'(x) = 0 \Rightarrow x = -2, x = 2$

x	$f(x)$
-6	3
-2	7
2	$7 - 2\pi$
5	$10 - 2\pi$

The absolute minimum value is $f(2) = 7 - 2\pi$.

(d) $f'''(-5) = \frac{2-0}{-6-(-2)} = -\frac{1}{2}$

2 : $\begin{cases} 1 : f'''(-5) \\ 1 : f'''(3) \text{ does not exist,} \\ \text{with explanation} \end{cases}$

$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} = 2$ and $\lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3} = -1$

$f'''(3)$ does not exist because

$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} \neq \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3}$