

Accelerated Pre-Calculus

February & March 2022


Unit 7 – Polar Graphs & Complex Numbers

Monday	Tuesday	Wednesday	Thursday	Friday
Feb 21 No School President's Day	22 7.01 Polar Coordinates <ul style="list-style-type: none"> • Plot points • Multiple representations HW: 7.01 Practice	23 7.02 Polar Coordinates <ul style="list-style-type: none"> • Convert btw Rectangular & Polar • Multiple representations • Distance Formula HW: 7.02 Practice	24 7.03 Polar Coordinate Review HW: Polar Review	25 7.04 Quiz- Polar Coordinates & Converting Points with Rectangular System
28 Early Release Day 7.05 Complex Numbers in Rectangular Form <ul style="list-style-type: none"> • Absolute Value • Modulus • Distance Between • Midpoint HW: 7.05 Practice	Mar 1 7.06 Adding & Subtracting Complex Coordinates Geometrically HW: 7.06 Practice	2 Check-In Quiz 7.07 Complex Numbers in Polar Form <ul style="list-style-type: none"> • Modulus and Argument HW: 7.07 Practice	3 7.08 Operations with Complex Numbers in Polar Form <ul style="list-style-type: none"> • Product • Quotient HW: 7.08 Practice	4 7.09 More Complex Number Operations <ul style="list-style-type: none"> • Power • Roots HW: 7.09 Practice
7 7.10 More Practice with Operations HW: 7.11 Review	8 7.11 Review HW: Study!	9 Test: Polar and Complex	10 TASK: Battleship - Star Wars Edition!	11 No School Teacher Work Day

Polar Coordinates, (r, θ) :

$$x = r \cos \theta$$

$$y = r \sin \theta$$



$$r = \sqrt{x^2 + y^2}$$

$$\theta = \tan^{-1}\left(\frac{y}{x}\right) \text{ if } x > 0$$

$$\text{or } \theta = \tan^{-1}\left(\frac{y}{x}\right) + \pi, \text{ if } x < 0$$

Distance between two points on the polar plane: $\sqrt{r_1^2 + r_2^2 - 2r_1r_2 \cos(\theta_2 - \theta_1)}$

Complex Numbers, Rectangular (Standard) form: $z = a + bi$

Absolute value (modulus): $|z| = \sqrt{a^2 + b^2}$

Distance between 2 complex numbers is the modulus of their difference: $|z_1 - z_2|$

Midpoint between 2 complex numbers is the average of the values: $\frac{z_1 + z_2}{2}$

Polar (Trigonometric) Form of a complex number: $z = r(\cos \theta + i \sin \theta)$ or $r \text{ cis } \theta$

Where $a = r \cos \theta$, $b = r \sin \theta$, $r = \sqrt{a^2 + b^2}$, and $\tan \theta = \frac{b}{a}$ (remember to add π if $a < 0$)

Multiplication of Complex Numbers

$$z_1 \cdot z_2 = r_1 r_2 [\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)]$$

Division of Complex Numbers

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} [\cos(\theta_1 - \theta_2) + i \sin(\theta_1 - \theta_2)], r_2 \neq 0$$

De Moivre's Theorem (Powers of a Complex Number)

$$z^n = [r(\cos \theta + i \sin \theta)]^n = r^n (\cos n\theta + i \sin n\theta)$$

 n th Roots of a Complex Number

$$\sqrt[n]{r} \left(\cos \frac{\theta + 2\pi k}{n} + i \sin \frac{\theta + 2\pi k}{n} \right), k = 0, 1, 2, \dots, n - 1$$