



1) Given the region below enclosed by $f(x) = \sqrt{x-2}$, $g(x) = 5-x$, and the x-axis.

a) Find the area of the below region. (Write the integral notation(s) as well as the numeric approximation rounded to 3 decimal places)



Method 2: Right-Left

$$\begin{array}{l|l} y = \sqrt{x-2} & y = 5-x \\ y^2 = x-2 & x = 5-y \\ y^2 + 2 = x & \end{array}$$




$$\text{Area} = \int_0^{1.303} (5-y - (y^2+2)) dy = \boxed{2.323}$$

Method 1: Top-Bottom, split into

2 regions: $\text{Area} = \int_2^{3.697} \sqrt{x-2} - 0 dx + \int_{3.697}^5 5-x - 0 dx = \boxed{2.323}$

b) Find the Volume of solid generated when the enclosed region is revolved about the line $x = 1$ (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)



$$R(y) = 5-y - (1) = 4-y$$


$$r(y) = y^2 + 2 - (1) = y^2 + 1$$

$$V = \pi \int_0^{1.303} [(4-y)^2 - (y^2+1)^2] dy$$

*Washer Method, Right-Left

$$\boxed{V = 11.265\pi \text{ units}^3}$$

c) The enclosed region is the base of a solid. The cross section of the solid taken perpendicular to the y-axis is an equilateral triangle. Find the volume of the given solid. (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)




$$\begin{aligned} \text{base} &= 5-y - (y^2+2) = 5-y-y^2-2 \\ &= 3-y^2-y \end{aligned}$$

$$\text{Area} = \frac{\sqrt{3}}{4} (\text{base})^2 \rightarrow \frac{\sqrt{3}}{4} (3-y^2-y)^2$$

$$V = \int_0^{1.303} \frac{\sqrt{3}}{4} (3-y^2-y)^2 dy = \boxed{2.225 \text{ units}^3}$$

2) Given the region below enclosed by $f(x) = \ln(4-x)$, the line $y=1$, and the y-axis.

a) Find the Volume of solid generated when the enclosed region is revolved about the line $y=1$ (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)




$$R(x) = \ln(4-x) - (1)$$

$$V = \pi \int_0^{1.282} [\ln(4-x) - 1]^2 dx$$

$$V = 0.069\pi \text{ units}^3$$

* Disc Method, Top-Bottom

b) Find the Volume of solid generated when the enclosed region is revolved about the line $y=2$ (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)



$$R(x) = 2 - (1) = 1$$


$$r(x) = 2 - (\ln(4-x))$$

$$V = \pi \int_0^{1.282} [1]^2 - [2 - \ln(4-x)]^2 dx$$

$$V = 0.457\pi \text{ units}^3$$

* washer Method, Right-Left

c) The enclosed region is the base of a solid. The cross section of the solid taken perpendicular to the x-axis is a rectangle whose height is twice the base. Find the volume of the given solid (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)



$$\text{base} = \ln(4-x) - (1)$$

$$\text{height} = 2[\ln(4-x) - 1]$$

$$\text{Area} = (\text{base})(\text{height})$$

$$\text{Area} = 2[\ln(4-x) - 1]^2$$


$$V = \int_0^{1.282} 2[\ln(4-x) - 1]^2 dx$$

$$V = 0.139 \text{ units}^3$$

Key

1)


Let f and g be the functions given by $f(x) = \frac{1}{4} + \sin(\pi x)$ and $g(x) = 4^{-x}$. Let R be the shaded region in the first quadrant enclosed by the y -axis and the graphs of f and g , and let S be the shaded region in the first quadrant enclosed by the graphs of f and g , as shown in the figure above.



a) Find the area of S


$$A = \int_{0.178}^1 \left(\frac{1}{4} + \sin(\pi x) - 4^{-x} \right) dx$$

$$A = 0.410 \text{ units}^2$$




b) Find the area of R

$$\text{Area} = \int_0^{0.178} \left(4^{-x} - \left(\frac{1}{4} + \sin(\pi x) \right) \right) dx = 0.0648 \text{ units}^2$$



c) Find the volume of the solid generated when S is revolved about the horizontal line $y = -1$.



*Washer Method


$$R(x) = \frac{1}{4} + \sin(\pi x) - (-1) = \frac{5}{4} + \sin(\pi x)$$

$$r(x) = 4^{-x} - (-1) = 4^{-x} + 1$$

$$V = \pi \int_{x_1}^{x_2} R(x)^2 - r(x)^2 dx$$

$$V = \pi \int_{0.178}^1 \left[\left(\frac{5}{4} + \sin(\pi x) \right)^2 - \left(4^{-x} + 1 \right)^2 \right] dx = 1.45/\pi \text{ units}^3$$

d) The enclosed region R is the base of a solid. The cross section of the solid taken parallel to the y -axis is a isosceles right triangle with leg on base. Find the volume of the given solid. (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)



$$\text{base} = 4^{-x} - \left(\frac{1}{4} + \sin(\pi x) \right)$$


$$\text{Area} = \frac{1}{2} (\text{base})^2 \rightarrow \frac{1}{2} \left[4^{-x} - \frac{1}{4} - \sin(\pi x) \right]^2$$

$$V = \int_{x_1}^{x_2} [\text{Area}] dx \rightarrow V = \int_0^{0.178} \frac{1}{2} \left[4^{-x} - \frac{1}{4} - \sin(\pi x) \right]^2 dx = 0.016 \text{ units}^3$$

2) Given the region below enclosed by $f(x) = \ln(4-x)$, the line $x=-1$, and the x-axis.

AOR
x=-1

a) Find the Volume of solid generated when the enclosed region is revolved about the line $x=-1$ (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)



* Disc Method

$$y = \ln(4-x) \quad \left| \quad e^y = 4-x \right.$$

$$e^y = e^{\ln(4-x)} \quad \left| \quad x = 4 - e^y \right.$$

$$e^y = (4-x)$$

(5)
x=-1
AOR


$$R(y) = 4 - e^y - (-1)$$

$$R(y) = 5 - e^y$$

$$V = \pi \int_{y_1}^{y_2} R(y)^2 dy$$

$$V = \pi \int_0^{1.61} [5 - e^y]^2 dy = \boxed{12.236\pi \text{ units}^3}$$

b) Find the Volume of solid generated when the enclosed region is revolved about the line $x=4$ (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)




$$R(y) = 4 - (-1) = 5$$

$$r(y) = 4 - (4 - e^y) = e^y$$

$$V = \pi \int_{y_1}^{y_2} R(y)^2 - r(y)^2 dy$$

$$V = \pi \int_0^{1.61} [5]^2 - [e^y]^2 dy = \boxed{28.236\pi \text{ units}^3}$$

c) The enclosed region is the base of a solid. The cross section of the solid taken parallel to the x-axis is a rectangle whose height is 4. Find the volume of the given solid. (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)



$$\text{base} = 4 - e^y - (-1) = 5 - e^y$$

$$\text{Area} = (\text{base})(\text{height})$$

$$\text{height} = 4$$

$$A = (5 - e^y)(4)$$



$$V = \int_{y_1}^{y_2} [\text{Area}] dy \rightarrow \int_0^{1.61} 4(5 - e^y) dy = \boxed{16.189 \text{ units}^3}$$

Key

1)


Let R be the region bounded by the graph of $y = e^{2x-x^2}$ and the horizontal line $y = 2$, and let S be the region bounded by the graph of $y = e^{2x-x^2}$ and the horizontal lines $y = 1$ and $y = 2$, as shown above.

- (a) Find the area of R .
- (b) Find the area of S .
- (c) Write, but do not evaluate, an integral expression that gives the volume of the solid generated when R is rotated about the horizontal line $y = 1$.



Top/bottom
 $y = e^{2x-x^2}$
 $y = 2$


$$\text{Area} = \int_{0.446}^{1.554} e^{2x-x^2} - 2 \, dx = \boxed{0.514}$$



Area of $S = \text{Area of } R+S - \text{Area of } R$

Area($R+S$) = $\int_{0.446}^{1.554} e^{2x-x^2} - 1 \, dx = 2.06016$
 (Top/bottom)

Area of $S = 2.06016 - 0.514 = \boxed{1.546}$



$R(x) = e^{2x-x^2} - 1$
 $r(x) = 2 - 1 = 1$

$$V = \pi \int_{x_1}^{x_2} R(x)^2 - r(x)^2 \, dx$$

$$V = \pi \int_{0.446}^{1.554} [e^{2x-x^2} - 1]^2 - [1]^2 \, dx$$


Washer Method

Top/bottom
 $y = e^{2x-x^2}$
 $y = 2$

2)


A baker is creating a birthday cake. The base of the cake is the region R in the first quadrant under the graph of $y = f(x)$ for

$0 \leq x \leq 30$, where $f(x) = 20 \sin\left(\frac{\pi x}{30}\right)$. Both x and y are measured in centimeters. The region R is shown in the figure above. The derivative of f is $f'(x) = \frac{2\pi}{3} \cos\left(\frac{\pi x}{30}\right)$.



(a) The region R is cut out of a 30-centimeter-by-20-centimeter rectangular sheet of cardboard, and the remaining cardboard is discarded. Find the area of the discarded cardboard.

(b) The cake is a solid with base R . Cross sections of the cake perpendicular to the x -axis are semicircles. If the baker uses 0.05 gram of unsweetened chocolate for each cubic centimeter of cake, how many grams of unsweetened chocolate will be in the cake?




$$\text{Area} = \text{Area of Box} - \text{Area under (Top-bottom) parabola}$$

$$y = 20 \sin\left(\frac{\pi x}{30}\right)$$

$$y = 0$$

$$= 30(20) - \int_0^{30} 20 \sin\left(\frac{\pi x}{30}\right) - 0 dx$$

$$= 600 - 381.972 = \boxed{218.028 \text{ cm}^2}$$



Top/bottom

$$y = 20 \sin\left(\frac{\pi x}{30}\right)$$

$$y = 0$$

$$\text{base} = 20 \sin\left(\frac{\pi x}{30}\right) - 0$$

$$\text{base} = 20 \sin\left(\frac{\pi x}{30}\right)$$

$$\text{Area} = \frac{\pi}{8} [\text{base}]^2$$

(semicircle)

$$= \frac{\pi}{8} \left[20 \sin\left(\frac{\pi x}{30}\right) \right]^2$$


$$\text{Volume} = \int_0^{30} \frac{\pi}{8} \left[20 \sin\left(\frac{\pi x}{30}\right) \right]^2 dx$$

$$V = 2356.194 \text{ cm}^3$$

$$\frac{0.05 \text{ grams}}{1 \text{ cm}^3} \cdot 2356.194 \text{ cm}^3$$


$$= \boxed{117.809 \text{ grams (of chocolate)}}$$

3) (Non-calculator)



Let R be the region in the first quadrant bounded by the graph of $y = 2\sqrt{x}$, the horizontal line $y = 6$, and the y -axis, as shown in the figure above.


- (a) Find the area of R .
- (b) Write, but do not evaluate, an integral expression that gives the volume of the solid generated when R is rotated about the horizontal line $y = 7$.
- (c) Region R is the base of a solid. For each y , where $0 \leq y \leq 6$, the cross section of the solid taken perpendicular to the y -axis is a rectangle whose height is 3 times the length of its base in region R . Write, but do not evaluate, an integral expression that gives the volume of the solid.

a) 
$$\begin{array}{l} \text{Top} - \text{bottom} \\ y = 2\sqrt{x} \\ y = 6 \end{array} \quad \left| \quad \text{Area} = \int_0^9 (6 - 2\sqrt{x}) dx \right.$$

$$= \int_0^9 (6 - 2x^{1/2}) dx = \left[6x - \frac{2x^{3/2}}{3/2} \right]_0^9$$

$$\left[6x - \frac{4}{3}x^{3/2} \right]_0^9 = 6(9) - \frac{4}{3}(9)^{3/2} - (0 - 0)$$

$$= 54 - \frac{4}{3}(27) = 54 - 4(9) = \boxed{18}$$



washer method

Top/bottom


$$y = 6$$

$$y = 2\sqrt{x}$$

$$R(x) = 7 - 2\sqrt{x}$$

$$r(x) = 7 - 6 = 1$$

$$V = \pi \int_0^9 [(7 - 2\sqrt{x})^2 - [1]^2] dx$$



$$y = 2\sqrt{x} \rightarrow \frac{y}{2} = \sqrt{x}$$

$$\left(\frac{y}{2}\right)^2 = x$$

$$\text{base} = \frac{y^2}{4} - 0 = \frac{y^2}{4}$$

$$\text{height} = 3(\text{base}) = 3\left(\frac{y^2}{4}\right)$$

$$\text{Area} = \text{base} \times \text{height}$$

$$= \left(\frac{y^2}{4}\right) \times 3\left(\frac{y^2}{4}\right) = \frac{3}{16}y^4$$



$$V = \int_0^6 \frac{3}{16}y^4 dy$$

4)

Let R be the shaded region bounded by the graph of $y = xe^{x^2}$, the line $y = -2x$, and the vertical line $x = 1$, as shown in the figure above.

(a) Find the area of R .

(b) Write, but do not evaluate, an integral expression that gives the volume of the solid generated when R is rotated about the horizontal line $y = -2$.



$$\text{Area} = \int_0^1 (xe^{x^2} - (-2x)) dx = \int_0^1 (xe^{x^2} + 2x) dx$$


$$\begin{aligned} u &= x^2 \\ \frac{du}{dx} &= 2x \\ dx &= \frac{du}{2x} \end{aligned} \quad \left| \int xe^u \frac{du}{2x} \right. \quad \left. \int \frac{1}{2} e^u du = \frac{1}{2} e^u \right.$$

u-substitution

$$\left. \left[\frac{1}{2} e^{x^2} + \frac{2x^2}{2} \right]_0^1 \right.$$

$$\frac{1}{2} e^1 + 1 - \left(\frac{1}{2} e^0 + 0^2 \right)$$

$$\frac{1}{2} e + 1 - \frac{1}{2} = \boxed{\frac{1}{2} e + \frac{1}{2}}$$



washer method

Top/bottom
 $y = xe^{x^2}$

$y = -2x$

$R(x) = xe^{x^2} - (-2)$

$r(x) = -2x - (-2)$
 $= -2x + 2$


$$V = \pi \int_{x_1}^{x_2} [R(x)]^2 - [r(x)]^2 dx$$

$$V = \pi \int_0^1 [xe^{x^2} + 2]^2 - [-2x + 2]^2 dx$$


Key

1. (Non-Calculator)

Let R be the region in the first quadrant enclosed by the graphs of $y = 2x$ and $y = x^2$, as shown in the figure above.



- (a) Find the area of R .
- (b) The region R is the base of a solid. For this solid, at each x the cross section perpendicular to the x -axis has area $A(x) = \sin\left(\frac{\pi}{2}x\right)$. Find the volume of the solid.
- (c) Another solid has the same base R . For this solid, the cross sections perpendicular to the y -axis are squares. Write, but do not evaluate, an integral expression for the volume of the solid.
- d) Find Volume of solid by rotating R about line $x = -1$




Top-bottom
 $y = 2x$
 $y = x^2$

Area = $\int_0^2 2x - x^2 dx$

$= \left[\frac{2x^2}{2} - \frac{x^3}{3} \right]_0^2 = 2^2 - \frac{2^3}{3} - (0 - 0)$
 $= 4 - \frac{8}{3} = \frac{12}{3} - \frac{8}{3} = \frac{4}{3}$

cross-section.



$V = \int [Area] dx$

$V = \int_0^2 \sin\left(\frac{\pi}{2}x\right) dx$

$u = \frac{\pi}{2}x$
 $\frac{du}{dx} = \frac{\pi}{2}$


$\pi dx = 2 du$
 $dx = \frac{2}{\pi} du$

$\int \sin u \cdot \frac{2}{\pi} du$

$\frac{2}{\pi} \int \sin u du$

$\left[-\frac{2}{\pi} \cos\left(\frac{\pi}{2}x\right) \right]_0^2$

$-\frac{2}{\pi} \cos(\pi) - \left(-\frac{2}{\pi} \cos(0)\right)$
 $= \frac{2}{\pi}(-1) + \frac{2}{\pi} = \frac{4}{\pi}$



base = $\sqrt{y} - \frac{y}{2}$


Area = [base]²
 square

Area = $\left[\sqrt{y} - \frac{y}{2}\right]^2$

$V = \int_0^4 \left[\sqrt{y} - \frac{y}{2}\right]^2 dy$

Right/Left

$x = \sqrt{y}$
 $x = \frac{y}{2}$



*washer method
 *Right/Left

$x = \sqrt{y}$
 $x = \frac{y}{2}$


$R(y) = \sqrt{y} - (-1)$
 $r(y) = \frac{y}{2} - (-1)$

$V = \pi \int_{y_1}^{y_2} [R(y)]^2 - [r(y)]^2 dy$

$V = \pi \int_0^4 \left[\sqrt{y} + 1\right]^2 - \left[\frac{y}{2} + 1\right]^2 dy$

2) (Non-Calculator)

Let $f(x) = 2x^2 - 6x + 4$ and $g(x) = 4\cos\left(\frac{\pi}{4}x\right)$. Let R be the region bounded by the graphs of f and g , as shown in the figure above.



- (a) Find the area of R .
- (b) Write, but do not evaluate, an integral expression that gives the volume of the solid generated when R is rotated about the horizontal line $y = 4$.
- (c) The region R is the base of a solid. For this solid, each cross section perpendicular to the x -axis is a square. Write, but do not evaluate, an integral expression that gives the volume of the solid.

a) $Area = \int_0^2 g(x) - f(x) dx$

$Area = \int_0^2 4\cos\left(\frac{\pi}{4}x\right) - (2x^2 - 6x + 4) dx$

$Area = \int_0^2 4\cos\left(\frac{\pi}{4}x\right) - 2x^2 + 6x - 4 dx$


$u = \frac{\pi}{4}x \quad dx = \frac{4}{\pi} du$
 $\frac{du}{dx} = \frac{\pi}{4} \quad 4 \int \cos u \cdot \frac{4}{\pi} du$
 $\pi dx = 4 du$

$\left[4 \cdot \frac{4}{\pi} \sin\left(\frac{\pi}{4}x\right) - \frac{2x^3}{3} + \frac{6x^2}{2} - 4x \right]_0^2$

$\frac{16}{\pi} \sin\left(\frac{\pi}{4} \cdot 2\right) - \frac{2(2)^3}{3} + \frac{6(2)^2}{2} - 4(2) - \left[\frac{16}{\pi} \sin(0) - 0 + 0 - 0 \right]$

$\frac{16}{\pi}(1) - \frac{16}{3} + \frac{24}{2} - 8$ or $\frac{16}{\pi} - \frac{4}{3}$

b) AOR: $y = 4$




* washer method
 * Top/bottom
 $y = 4\cos\left(\frac{\pi}{4}x\right)$
 $y = 2x^2 - 6x + 4$

$R(x) = 4 - (2x^2 - 6x + 4) = 4 - 2x^2 + 6x - 4 = -2x^2 + 6x$
 $r(x) = 4 - 4\cos\left(\frac{\pi}{4}x\right)$

$V = \pi \int_0^2 \left[-2x^2 + 6x \right]^2 - \left[4 - 4\cos\left(\frac{\pi}{4}x\right) \right]^2 dx$

$V = \int_0^2 \left[4\cos\left(\frac{\pi}{4}x\right) - 2x^2 + 6x - 4 \right]^2 dx$

c)




Top/bottom
 $y = 4\cos\left(\frac{\pi}{4}x\right)$
 $y = 2x^2 - 6x + 4$

base = $4\cos\left(\frac{\pi}{4}x\right) - (2x^2 - 6x + 4)$
 Area square = $[base]^2$

Key

1) Given the region below enclosed by $f(x) = \ln(x-3)$, the line $y = 7 - \frac{1}{4}x$, and the x -axis.



Method 1: (Top-Bottom) * 2 separate integrals *

$$\text{Area} = \int_4^{17.346} \ln(x-3) - 0 \, dx + \int_{17.346}^{28} 7 - \frac{1}{4}x - 0 \, dx$$

Method 2: (Right-Left)

$$y = \ln(x-3) \quad | \quad y = 7 - \frac{1}{4}x$$


$$e^y = x - 3 \quad | \quad (y - 7 = -\frac{1}{4}x) \cdot 4$$

$$3 + e^y = x \quad | \quad 28 - 4y = x$$

$$\text{Area} = 24.864 + 14.188 = \boxed{39.052 \text{ units}^2} \quad \text{Area} = \int_0^{2.663} 28 - 4y - (3 + e^y) \, dy = \boxed{39.052 \text{ units}^2}$$

2) Given the region below enclosed by $f(x) = \ln(x+6)$, the line $y = -3$, and $x = 5$.

a) Find the Volume of solid generated when the enclosed region is revolved about the line $y = -4$ (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)



* Washer Method (Top-Bottom)


$$R(x) = \ln(x+6) - (-4) = \ln(x+6) + 4$$

$$r(x) = -3 - (-4) = 1$$

$$V = \pi \int_{x_1}^{x_2} R(x)^2 - r(x)^2 \, dx$$

$$V = \pi \int_{-5.95}^5 [\ln(x+6) + 4]^2 - [1]^2 \, dx = \boxed{320.510\pi \text{ units}^3}$$

b) Find the Volume of solid generated when the enclosed region is revolved about the line $x = 5$ (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)



* find upper bound:

$$y(x) = \ln(x+6)$$

$$y(5) = \ln(5+6) = \ln 11 = 2.398$$

$$V = \pi \int_{y_1}^{y_2} [R(y)]^2 \, dy$$

* Disc Method (Right-Left)

(5 AOR $x=5$)


$$R(y) = 5 - (e^y - 6)$$

$$R(y) = 11 - e^y$$

$$V = \pi \int_{-3}^{2.398} [11 - e^y]^2 \, dy = \boxed{472.739\pi \text{ units}^3}$$

3) Given the region below enclosed by $f(x) = \sqrt{x+6}$, the $g(x) = e^x + 1$

a) Find the Volume of solid generated when the enclosed region is revolved about the line $x = -6$ (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)



$x = y^2 - 6$ (0.429, 2.536)
 $y = \sqrt{x+6}$
 $(y)^2 = (\sqrt{x+6})^2$
 $y^2 = x+6$
 $y^2 - 6 = x$

$y = e^x + 1$
 $y - 1 = e^x$
 $\ln(y-1) = \ln e^x$
 $\ln(y-1) = x \ln e$
 $\ln(y-1) = x$


$V = \pi \int_{y_1}^{y_2} (R(y)^2 - r(y)^2) dy$
 $V = 26.032\pi$
 units³

$R(y) = \ln(y-1) - (-6) = \ln(y-1) + 6$
 $r(y) = y^2 - 6 - (-6) = y^2$

$V = \pi \int_{1.007}^{2.536} [(\ln(y-1) + 6)^2 - (y^2)^2] dy$

(S) AOR $x = -6$
 * Washer Method (Right-Left)

b) The enclosed region is the base of a solid. The cross section of the solid taken parallel to the y-axis is an isosceles right triangle with hypotenuse on base. Find the volume of the given solid. (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)




$V = \int_{x_1}^{x_2} [\text{Area}] dx$
 $V = \int_{-4.986}^{0.429} \frac{1}{4} [\sqrt{x+6} - e^x - 1]^2 dx$
 $V = 0.581$
 units³

base = $\sqrt{x+6} - (e^x + 1)$
 base = $\sqrt{x+6} - e^x - 1$

$\text{Area} = \frac{1}{4} (\text{base})^2$
 $\text{Area} = \frac{1}{4} (\sqrt{x+6} - e^x - 1)^2$

4) Given the region below enclosed by $f(x) = -x^2 + 4$ and $g(x) = -\frac{1}{2}x + 2$

Find the Volume of solid generated when the enclosed region is revolved about the line $y = 4$ (Write the integral notation as well as the numeric approximation rounded to 3 decimal places)



AOR $y = 4$
 * Washer Method (Top-Bottom)

$R(x) = 4 - (-\frac{1}{2}x + 2) = 4 + \frac{1}{2}x - 2 = 2 + \frac{1}{2}x$
 $r(x) = 4 - (-x^2 + 4) = 4 + x^2 - 4 = x^2$

$V = \pi \int_{x_1}^{x_2} (R(x)^2 - r(x)^2) dx$
 $V = \pi \int_{-1.186}^{1.686} [2 + \frac{1}{2}x]^2 - [x^2]^2 dx = 10.268\pi$
 units³